arXiv:1612.08572v1 [math.PR] 27 Dec 2016

Uniform infinite half-planar quadrangulations with skewness

Erich Baur*, Loïc Richier†
ENS Lyon

December 28, 2016

Abstract

We introduce a one-parameter family of random infinite quadrangulations of the half-plane, which we call the *uniform infinite half-planar quadrangulations with skewness* (UIHPQ_p for short, with $p \in [0, 1/2]$ measuring the skewness). They interpolate between Kesten's tree corresponding to $p = 0$ and the usual UIHPQ with a general boundary corresponding to $p = 1/2$. As we make precise, these models arise as local limits of uniform quadrangulations with a boundary when their volume and perimeter grow in a properly fine-tuned way, and they represent all local limits of (sub)critical Boltzmann quadrangulations whose perimeter tend to infinity. Our main result shows that the family $(\mathsf{UIHPQ}_p)_p$ approximates the Brownian half-planes BHP_θ, $\theta \geqslant 0$, recently introduced in [8]. For $p < 1/2$, we give a description of the UIHPQ_p in terms of a looptree associated to a critical two-type Galton-Watson tree conditioned to survive.

Key words: Uniform infinite half-planar quadrangulation, Brownian half-plane, Kesten's tree, multi-type Galton-Watson tree, looptree, Boltzmann map.
Subject Classification: 05C80; 05C81; 05C05; 60J80; 60F05.

Contents

1 Introduction **2**
1.1 Overview 2
1.2 Some standard notation and definitions 5

*erich.baur@ens-lyon.fr
†loic.richier@ens-lyon.fr

Acknowledgment of support. The research of EB was supported by the Swiss National Science Foundation grant P300P2_161011, and performed within the framework of the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR). EB and LR thank the Institute for Mathematical Research (FIM) of ETH Zurich for hospitality, where parts of this work were completed.

2 Statements of the main results **8**
2.1 Local limits 8
2.2 Scaling limits 9
2.3 Tree structure 11

3 Random half-planes and trees **13**
3.1 The Brownian half-planes BHP_θ 13
3.2 Random trees and some of their properties 15

4 Construction of the UIHPQ_p **18**
4.1 The encoding objects 18
4.2 The Bouttier-Di Francesco-Guitter mapping 20
4.3 Definition of the UIHPQ_p 22

5 Proofs of the limit results **24**
5.1 The UIHPQ_p as a local limit of uniform quadrangulations 24
5.2 The UIHPQ_p as a local limit of Boltzmann quadrangulations 31
5.3 The BHP_θ as a local scaling limit of the UIHPQ_p's 33
5.4 The ICRT as a local scaling limit of the UIHPQ_p's 37

6 Proofs of the structural properties **42**
6.1 The branching structure behind the UIHPQ_p 42
6.2 Recurrence of simple random walk 46

Literature **47**

1 Introduction

1.1 Overview

The purpose of this paper is to introduce and study a one-parameter family of random infinite quadrangulations of the half-plane, which we denote by $(\mathsf{UIHPQ}_p)_{0\leqslant p\leqslant 1/2}$ and call the *uniform infinite half-planar quadrangulations with skewness*. Two members play a particular role: The choice $p = 0$ corresponds to Kesten's tree, cf. Proposition 1 below, whereas the choice $p = 1/2$ corresponds to the standard uniform infinite half-planar quadrangulation UIHPQ with a general boundary.

Kesten's tree [30] is a random infinite planar tree, which we may view as a degenerate quadrangulation with an infinite boundary, but no inner faces. It arises as the local limit of critical Galton-Watson trees conditioned to survive. The standard $\mathsf{UIHPQ}(= \mathsf{UIHPQ}_{1/2})$ forms the half-planar analog of the uniform infinite planar quadrangulation introduced by Krikun [31], after the seminal work of Angel and Schramm [7] on triangulations of the plane. Curien and Miermont [25] showed that the UIHPQ arises as a local limit of uniformly chosen quadrangulations of the two-sphere with n inner faces and a boundary of size 2σ, upon letting first $n \to \infty$ and then $\sigma \to \infty$ (see Angel [3] for the case of triangulations with a simple boundary).

We will define each UIHPQ_p in Section 4 by means of an extension of the Bouttier-Di Francesco-Guitter mapping to infinite quadrangulations with a boundary. In the first part of this paper, we will discuss various local limits and scaling limits which involve the family $(\mathsf{UIHPQ}_p)_p$. More precisely, in Theorem 1, we will see that each UIHPQ_p appears as a local limit as n tends to infinity of uniform quadrangulations $Q_n^{\sigma_n}$ with n inner faces and a boundary of size $2\sigma_n$, for an appropriate choice of $\sigma_n = \sigma_n(p) \to \infty$. In Proposition 2, we argue that the family $(\mathsf{UIHPQ}_p)_p$ consists precisely of the infinite quadrangulations with a boundary which are obtained as local limits $\sigma \to \infty$ of subcritical Boltzmann quadrangulations with a boundary of size 2σ. This result will prove helpful in our description of the UIHPQ_p given in Theorem 4.

We will then turn to distributional scaling limits of the family $(\mathsf{UIHPQ}_p)_p$ in the so-called *local Gromov-Hausdorff topology*. In Theorems 2 and 3, we will clarify the connection between the (discrete) quadrangulations UIHPQ_p and the family $(\mathsf{BHP}_\theta)_{\theta \geqslant 0}$ of Brownian half-spaces with skewness θ introduced in [8]. More specifically, upon rescaling the graph distance by a factor $a_n^{-1} \to 0$, we prove that each BHP_θ is the distributional limit of the rescaled spaces $a_n^{-1} \cdot \mathsf{UIHPQ}_{p_n}$, if $p_n = p_n(\theta, a_n)$ is adjusted in the right manner (Theorem 2). In our setting, convergence in the local Gromov-Hausdorff sense amounts to show convergence of rescaled metric balls around the roots of a fixed but arbitrarily large radius in the usual Gromov-Hausdorff topology; see Section 1.2.7.

In [8], a classification of all possible non-compact scaling limits of pointed uniform random quadrangulations with a boundary $(V(Q_n^{\sigma_n}), a_n^{-1} d_{\mathrm{gr}}, \rho_n)$ has been given, depending on the asymptotic behavior of the boundary size $2\sigma_n$ and on the choice of the scaling factor $a_n \to \infty$ (in the local Gromov-Hausdorff topology, with the distinguished point ρ_n lying on the boundary). In this paper, we address the boundary regime corresponding to the portion $x \geqslant 1$ of the $y = 0$ axis in Figure 1 (in hashed marks), which was left untouched in [8]. As we show, it corresponds to a regime of unrescaled local limits, namely the family $(\mathsf{UIHPQ}_p)_p$.

We finally give a branching characterization of the UIHPQ_p when $p < 1/2$. For that purpose, we will adapt the concept of discrete random looptrees introduced by Curien and Kortchemski [22]. We will see that the UIHPQ_p admits a representation in terms of a looptree associated to a two-type version of Kesten's infinite tree. Informally, we will replace each vertex u at odd height in Kesten's tree by a cycle of length $\deg(u)$, which connects the vertices incident to u. Here, $\deg(u)$ stands for the degree (i.e., the number of neighbors) of u in the tree. We then fill in the cycles of the looptree with a collection of independent quadrangulations with a simple boundary, which are drawn according to a subcritical Boltzmann law. As we show in Theorem 4, the space constructed in this way has the law of the UIHPQ_p. Discrete looptrees and their scaling limits have found various applications in the study of large-scale properties of random planar maps, for instance in the description of the boundary of percolation clusters on the uniform infinite planar triangulation; see the work [23], which served as the main inspiration for our characterization of the UIHPQ_p. From our description, we immediately infer that simple random walk is recurrent on the UIHPQ_p for $p < 1/2$.

It is well-known that the standard UIHPQ with a simple boundary satisfies the so-called *spatial Markov property*, which allows, in particular, the use of peeling techniques. In [5], Angel and Ray classified all triangulations (without self-loops) of the half-plane satisfying the spatial Markov property and translation invariance. They form a one-parameter



Figure 1: In [8], all possible limits for the rescaled spaces $(V(Q_n^{\sigma_n}), a_n^{-1} d_{\mathrm{gr}}, \rho_n)$ are discussed. The x-axis represents the limit values for the logarithm of the boundary length $\log(\sigma_n)/\log(n)$ in units of $\log(n)$, and the y-axis corresponds to the limit of the logarithm of the scaling factor $\log(a_n)/\log(n)$ in units of $\log(n)$. The focus of this paper lies on the hashed region.

family $(\mathbb{H}_\alpha)_\alpha$ parametrized by $\alpha \in [0, 1)$. The parameter $\alpha = 2/3$ corresponds to the standard UIHPT with a simple boundary, the triangular equivalent of the UIHPQ with a simple boundary. When $\alpha > 2/3$ (the supercritical case), $\mathbb{H}_\alpha$ is of hyperbolic nature and exhibits an exponential volume growth. On the contrary, when $\alpha < 2/3$ (the subcritical case), it has a tree-like structure. We believe that the family $(\mathsf{UIHPQ}_p)_p$ is a quadrangular equivalent to the triangulations in the subcritical phase of [5]. Note that contrary to the UIHPQ_p, the spaces $\mathbb{H}_\alpha$ for $\alpha < 2/3$ have a half-plane topology, due to the conditioning to have a simple boundary. However, there exists almost surely infinitely many cut-edges connecting the left and right boundaries; see [37, Proposition 4.11]. This should be seen as an equivalent to the branching structure formulated in Theorem 4 below. Our methods in this paper are different from [5, 37] as we do not use peeling techniques.

In [21], Curien studied full-plane analogs of the family $(\mathbb{H}_\alpha)_\alpha$. With similar (peeling) techniques, he constructed a (unique) one-parameter family of random infinite planar triangulations indexed by $\kappa \in (0, 2/27]$, which satisfy a slightly adapted spatial Markov property. The critical case $\kappa = 2/27$ corresponds to the standard UIPT with a simple boundary of Angel and Schramm [7]. The regime $\kappa \in (0, 2/27)$ parallels the supercritical (or hyperbolic) phase $\alpha > 2/3$ of [5], whereas it is shown that there is no subcritical phase. Recently, a near-critical scaling limit of hyperbolic nature called the hyperbolic Brownian half-plane has been studied by Budzinski [17]. It is obtained from rescaling the triangulations of Curien [21] and letting $\kappa \to 2/27$ at the right speed. Theorem 1 of [17] bears some structural similarities with our Theorem 2 below, although it concerns a different regime.

Structure of the paper

The rest of this paper is structured as follows. In the following section, we introduce some (standard) concepts and notation around quadrangulations, which will be used throughout

this text. Moreover, we recapitulate the local topology and the local Gromov-Hausdorff topology. In Section 2, we state our main results, which concern local limits, scaling limits, and structural properties of the family $(\mathsf{UIHPQ}_p)_p$. Section 3 reviews the definition of the family of Brownian half-planes $(\mathsf{BHP}_\theta)_\theta$, and of various random trees, which are used both to describe the distributional limits of the family $(\mathsf{UIHPQ}_p)_p$ as well as their branching structure.

In Section 4, we construct the UIHPQ_p. We first explain the Bouttier-Di Francesco-Guitter encoding of quadrangulations with a boundary and then define the UIHPQ_p in terms of the encoding objects. We are then in position to prove our limit statements; see Section 5. In the final Section 6, we prove our main result characterizing the tree-like structure of the UIHPQ_p when $p < 1/2$, as well as recurrence of simple random walk.

1.2 Some standard notation and definitions

1.2.1 Notation

We write

$$\mathbb{N} = \{1, 2, \ldots\}, \quad \mathbb{N}_0 = \mathbb{Z}_{\geqslant 0} = \mathbb{N} \cup \{0\}, \quad \mathbb{Z}_{<0} = \{-1, -2, \ldots\}.$$

For two sequences $(a_n)_n$, $(b_n)_n \subset \mathbb{N}$, we write $a_n \ll b_n$ or $b_n \gg a_n$ if $a_n/b_n \to 0$ as $n \to \infty$. Given two measurable subsets $U, V \subset \mathbb{R}$, we denote by $\mathcal{C}(U, V)$ the space of continuous functions from U to V, equipped with the usual compact-open topology, i.e., uniform convergence on compact subsets. We write $\|\nu\|_{\mathrm{TV}}$ for the total variation norm of a probability measure ν.

As a general notational rule for this paper, if we drop p from the notation, we work with the case $p = 1/2$. For example, we write UIHPQ (and not $\mathsf{UIHPQ}_{1/2}$) for the standard uniform infinite half-planar quadrangulation.

1.2.2 Planar maps

By a *planar map* we mean, as usual, an equivalence class of a proper embedding of a finite connected graph in the two-sphere, where two embeddings are declared to be equivalent if they differ only by an orientation-preserving homeomorphism of the sphere. Loops and multiple edges are allowed. Our planar maps will be rooted, meaning that we distinguish an oriented edge called the *root edge*. Its origin is the root vertex of the map. The faces of a planar map are formed by the components of the complement of the union of its edges.

1.2.3 Quadrangulations with a boundary

A *quadrangulation with a boundary* is a finite planar map q, whose faces are quadrangles except possibly one face called the *outer face*, which may an have arbitrary even degree. The edges incident to the outer face form the *boundary* $\partial\mathsf{q}$ of q, and their number $\#\partial\mathsf{q}$ (counted with multiplicity) is the size or *perimeter* of the boundary. In general, we do not assume that the boundary edges form a simple curve. We will root the map by selecting an oriented edge of the boundary, such that the outer face lies to its right. The *size* of q is given by the number of its *inner faces*, i.e., all the faces different from the outer face.

We write $\mathcal{Q}_n^\sigma$ for the (finite) set of all rooted quadrangulations with n inner faces and a boundary of size 2σ, $\sigma \in \mathbb{N}_0$. By convention, $\mathcal{Q}_0^0 = \{\dagger\}$ consists of the unique vertex map.

More generally, $\mathcal{Q}_f$ will denote the set of all finite rooted quadrangulations with a boundary, and $\mathcal{Q}_f^\sigma \subset \mathcal{Q}_f$ the set of all finite rooted quadrangulations with 2σ boundary edges, for $\sigma \in \mathbb{N}_0$.

Similarly, we let $\widehat{\mathcal{Q}}_f$ be the set of all finite rooted quadrangulations with a simple boundary, meaning that the edges of their outer face form a cycle without self-intersection. We denote by $\widehat{\mathcal{Q}}_f^\sigma \subset \widehat{\mathcal{Q}}_f$ the subset of finite rooted quadrangulations with a simple boundary of size 2σ. Note that $\mathcal{Q}_0^1$ consists of the map having one oriented edge and thus a simple boundary.

1.2.4 Uniform quadrangulations with a boundary

Throughout this text, we write Q_n^σ for a quadrangulation chosen uniformly at random in $\mathcal{Q}_n^\sigma$. We denote by ρ_n the root vertex of Q_n^σ, i.e., the origin of the root edge. By equipping the set of vertices $V(Q_n^\sigma)$ with the graph distance d_{gr}, we view the triplet $(V(Q_n^\sigma), d_{\mathrm{gr}}, \rho_n)$ as a random rooted metric space.

1.2.5 Boltzmann quadrangulations with a boundary

We will also work with various Boltzmann measures. For a finite rooted quadrangulation $\mathsf{q} \in \mathcal{Q}_f$, we write $\mathrm{F}(\mathsf{q})$ for the set of inner faces of q. Given non-negative weights g per inner face and $\sqrt{z}$ per boundary edge, we let

$$F(g,z) = \sum_{\mathsf{q}\in\mathcal{Q}_f} g^{\#\mathrm{F}(\mathsf{q})} z^{\#\partial\mathsf{q}/2}.$$

When this partition function is finite, we may define the associated Boltzmann distribution

$$\mathbb{P}_{g,z}(\mathsf{q}) = \frac{g^{\#\mathrm{F}(\mathsf{q})} z^{\#\partial\mathsf{q}/2}}{F(g,z)}, \quad \mathsf{q} \in \mathcal{Q}_f.$$

The statement of Proposition 2 below deals with Boltzmann-distributed quadrangulations of a fixed boundary size 2σ, for $\sigma \in \mathbb{N}_0$. In this case, the associated partition function and Boltzmann distribution read

$$F_\sigma(g) = \sum_{\mathsf{q}\in\mathcal{Q}_f^\sigma} g^{\#\mathrm{F}(\mathsf{q})}, \quad \mathbb{P}_g^\sigma(\mathsf{q}) = \frac{g^{\#\mathrm{F}(\mathsf{q})}}{F_\sigma(g)}, \quad \mathsf{q} \in \mathcal{Q}_f^\sigma,$$

whenever $g \geqslant 0$ is such that $F^\sigma(g)$ is finite. The Boltzmann distribution $\mathbb{P}_g^\sigma$ is related to $\mathbb{P}_{g,z}$ by conditioning the latter with respect to the boundary length, i.e., $\mathbb{P}_g^\sigma(\mathsf{q}) = \mathbb{P}_{g,z}(\mathsf{q} \mid \mathcal{Q}_f^\sigma)$.

When studying quadrangulations with a simple boundary, the partition functions are

$$\widehat{F}(g,z) = \sum_{\mathsf{q}\in\widehat{\mathcal{Q}}_f} g^{\#\mathrm{F}(\mathsf{q})} z^{\#\partial\mathsf{q}/2}, \quad \widehat{F}_\sigma(g) = \sum_{\mathsf{q}\in\widehat{\mathcal{Q}}_f^\sigma} g^{\#\mathrm{F}(\mathsf{q})},$$

and the Boltzmann distributions take the form

$$\widehat{\mathbb{P}}_{g,z}(\mathsf{q}) = \frac{g^{\#\mathrm{F}(\mathsf{q})}}{\widehat{F}_\sigma(g,z)}, \quad \mathsf{q} \in \widehat{\mathcal{Q}}_f, \quad \widehat{\mathbb{P}}_g^\sigma(\mathsf{q}) = \frac{g^{\#\mathrm{F}(\mathsf{q})}}{\widehat{F}_\sigma(g)}, \quad \mathsf{q} \in \widehat{\mathcal{Q}}_f^\sigma.$$

Remark 1. In the notation of [15], the generating function F is denoted W_0, while $\widehat{F}$ is denoted $\tilde{W}_0$. The index zero stands for the distance between the origin of the root edge and the marked vertex, so that these generating functions count unpointed quadrangulations.

1.2.6 Local topology

Our unrescaled limit results hold with respect to the *local topology* first studied by Benjamini and Schramm [10]: For two rooted planar maps m and m', the local distance between m and m' is

$$d_{\mathrm{map}}(\mathsf{m},\mathsf{m}') = (1+\sup\{r\geqslant 0 : \mathrm{Ball}_r(\mathsf{m}) = \mathrm{Ball}_r(\mathsf{m}')\})^{-1},$$

where $\mathrm{Ball}_r(\mathsf{m})$ denotes the combinatorial ball of radius r around the root ρ of m, i.e., the submap of m consisting of all the vertices v of m with $d_{\mathrm{gr}}(\rho,v)\leqslant r$ and all the edges of m between such vertices. The set $\mathcal{Q}_f$ of all finite rooted quadrangulations with a boundary is not complete for the distance d_{map}; we have to add infinite quadrangulations. We shall write $\mathcal{Q}$ for the completion of $\mathcal{Q}_f$ with respect to d_{map}. The UIHPQ_p will be defined as a random element in $\mathcal{Q}$.

1.2.7 Around the Gromov-Hausdorff metric

The pointed Gromov-Hausdorff distance measures the distance between (pointed) compact metric spaces, where the latter are viewed up to isometries. More specifically, given two elements $\mathbf{E}=(E,d,\rho)$ and $\mathbf{E}'=(E',d',\rho')$ in the space $\mathbb{K}$ of isometry classes of pointed compact metric spaces, their Gromov-Hausdorff distance is defined as

$$d_{\mathrm{GH}}(\mathbf{E},\mathbf{E}') = \inf\left\{d_{\mathrm{H}}(\varphi(E),\varphi'(E))\vee\delta(\varphi(\rho),\varphi'(\rho'))\right\},$$

where the infimum is taken over all isometric embeddings $\varphi : E\to F$ and $\varphi' : E'\to F$ of E and E' into the same metric space (F,δ), and d_{H} is the usual Hausdorff distance between compacts of F. The space $(\mathbb{K},d_{\mathrm{GH}})$ is complete and separable.

Our results on scaling limits involve non-compact pointed metric spaces and hold in the so-called *local Gromov-Hausdorff sense*, which we briefly recall next. Given a pointed complete and locally compact length space $\mathbf{E}$ and a sequence $(\mathbf{E}_n)_n$ of such spaces, $(\mathbf{E}_n)_n$ converges in the local Gromov-Hausdorff sense to $\mathbf{E}$ if for every $r\geqslant 0$,

$$d_{\mathrm{GH}}(B_r(\mathbf{E}_n),B_r(\mathbf{E}))\to 0 \quad \text{as } n\to\infty.$$

Here and in what follows, given a pointed metric space $\mathbf{F}=(F,d,\rho)$, $B_r(\mathbf{F})=\{x\in F : d(x,\rho)\leqslant r\}$ denotes the closed ball of radius r around ρ, viewed as a subspace of $\mathbf{F}$ equipped with the metric structure inherited from $\mathbf{F}$. For $\lambda>0$, $\lambda\cdot\mathbf{F}$ stands for the rescaled pointed metric space $(F,\lambda d,\rho)$, so that in particular $\lambda\cdot B_r(\mathbf{F}) = B_{\lambda r}(\lambda\cdot\mathbf{F})$.

As a discrete map, the UIHPQ_p is not a length space in the sense of [18]. However, by identifying each edge with a copy of the unit interval $[0,1]$ (and by extending the metric isometrically), one obtains a complete locally compact length space (pointed at the root vertex). By construction, balls of the same radius and around the same points in the UIHPQ_p and in the approximating length space are at Gromov-Hausdorff distance at most 1 from each other. Therefore, local Gromov-Hausdorff convergence for the (rescaled) UIHPQ_p, see Theorems 2 and 3 below, follows indeed from the convergence of balls as stated above.

2 Statements of the main results

2.1 Local limits

Our first result states that each member of the family $(\mathsf{UIHPQ}_p)_{0\leqslant p\leqslant 1/2}$ can be seen as a local limit $n\to\infty$ of uniform quadrangulations of with n inner faces and a boundary of size $2\sigma_n$, provided $\sigma_n=\sigma_n(p)$ is chosen in the right manner.

Theorem 1. *Fix $0\leqslant p\leqslant 1/2$, and let $(\sigma_n, n\in\mathbb{N})$ be a sequence of positive integers satisfying*

$$\sigma_n=\frac{1-2p}{p}n+o(n)\quad \text{if } 0<p\leqslant 1/2,\quad \text{and}\quad \sigma_n\gg n\quad \text{if } p=0.$$

For every $n\in\mathbb{N}$, let $Q_n^{\sigma_n}$ be uniformly distributed in $\mathcal{Q}_n^{\sigma_n}$. Then we have the local convergence for the metric d_{map} as $n\to\infty$,

$$Q_n^{\sigma_n}\xrightarrow{(d)}\mathsf{UIHPQ}_p.$$

In fact, we will prove a stronger result than mere local convergence: We will establish an isometry of balls of growing radii around the roots, where the maximal growth rate of the radii is given by $\xi_n=\min\{n^{1/4},\sqrt{n/\gamma_n}\}$, for $\gamma_n=\max\{\sigma_n-\frac{1-2p}{p}n,1\}$. We defer to Proposition 4 for the exact statement. The case $p=1/2$ corresponding to the regime $\sigma_n=o(n)$ is already covered by [8, Proposition 3.11] and is only included for completeness.

The convergence in the case $p=0$ with $\sigma_n\gg n$ is somewhat simpler. However, it is *a priori* not obvious that the UIHPQ_0 as defined in Section 4 is actually Kesten's tree (see Section 3.2.3 for a definition of the latter).

Proposition 1. *The space UIHPQ_0 has the law of Kesten's tree $\mathcal{T}_\infty$ associated to the critical geometric probability distribution $(\mu_{1/2}(k),k\in\mathbb{N}_0)$ given by $\mu_{1/2}(k)=2^{-(k+1)}$.*

Interestingly, the fact that the UIHPQ_0 is Kesten's tree can also be derived as a special case from Theorem 4 below; see Remark 5. We prefer, however, to give a direct proof of the proposition based on our construction of the UIHPQ_0.

The UIHPQ_p for $0\leqslant p\leqslant 1/2$ is also obtained as a local limit of Boltzmann quadrangulations with growing boundary size. This result will be important to describe the tree-like structure of the UIHPQ_p when $p<1/2$. More specifically, the family $(\mathsf{UIHPQ}_p)_p$ is precisely given by the collection of all local limits $\sigma\to\infty$ of Boltzmann quadrangulations with a boundary of size 2σ and weight $g\leqslant g_c=1/12$ per inner face. The value $g_c=1/12$ is critical (see [15, Section 4.1]) and corresponds to the choice $p=1/2$.

Proposition 2. *Fix $0\leqslant p\leqslant 1/2$, and set $g_p=p(1-p)/3$. For every $\sigma\in\mathbb{N}_0$, let $Q_\sigma(p)$ be a random rooted quadrangulation distributed according to the Boltzmann measure $\mathbb{P}_{g_p}^\sigma$. Then we have the local convergence for the metric d_{map} as $\sigma\to\infty$,*

$$Q_\sigma(p)\xrightarrow{(d)}\mathsf{UIHPQ}_p.$$

Remark 2. For $p=1/2$, the above proposition states convergence of critical Boltzmann quadrangulations with a boundary towards the UIHPQ, as it was already proved in [20, Theorem 7] by means of peeling techniques. In view of the above proposition, it is moreover

implicit from the same theorem that an infinite random map with the law of the UIHPQ_p does exist. For the case of half-planar triangulations (with a simple boundary), see [3]. When $p = 0$, there is no inner quadrangle almost surely and $Q^\sigma(0)$ is a uniform tree with σ edges (i.e., a Galton-Watson tree with geometric offspring law conditioned to have σ edges), which converges locally towards Kesten's tree; see, for example, [28, Theorem 7.1].

Remark 3. Let us write $\mathcal{M}(\mathcal{Q})$ for the set of probability measures on the completion $\mathcal{Q}$, and equip it with the usual weak topology. Then it is easily seen by our methods that the mapping $[0, 1/2] \ni p \mapsto \mathrm{Law}(\mathsf{UIHPQ}_p) \in \mathcal{M}(\mathcal{Q})$ is continuous.

2.2 Scaling limits

Our next results address scaling limits of the family $(\mathsf{UIHPQ}_p)_p$. In [8], a one-parameter family of (non-compact) random rooted metric spaces called the *Brownian half-planes* BHP_θ with skewness $\theta \geqslant 0$ was introduced. See Section 3.1 for a quick reminder. The Brownian half-plane BHP_0 corresponding to the choice $\theta = 0$ forms the half-planar analog of the Brownian plane introduced in [24] and arises from zooming-out the UIHPQ around the root vertex; see [8, Theorem 3.6], and [27, Theorem 1.10]). Here, we will see more generally that the family $(\mathsf{UIHPQ}_p)_p$ approximates the space BHP_θ for each $\theta \geqslant 0$ in the local Gromov-Hausdorff sense, provided p is appropriately fine-tuned (depending on θ).

Theorem 2. *Let $\theta \geqslant 0$. Let $(a_n, n \in \mathbb{N})$ be a sequence of positive reals with $a_n \to \infty$ as $n \to \infty$. Let $(p_n, n \in \mathbb{N}) \subset [0, 1/2]$ be a sequence satisfying*

$$p_n = p_n(\theta, a_n) = \frac{1}{2}\left(1 - \frac{2\theta}{3a_n^2}\right) + o\left(a_n^{-2}\right).$$

Then, in the sense of the local Gromov-Hausdorff topology as $n \to \infty$,

$$a_n^{-1} \cdot \mathsf{UIHPQ}_{p_n} \xrightarrow{(d)} \mathsf{BHP}_\theta.$$

The space BHP_θ satisfies the scaling property $\lambda \cdot \mathsf{BHP}_\theta =_d \mathsf{BHP}_{\theta/\lambda^2}$. It was shown in Remark 3.19 of [8] that Aldous' infinite continuum random tree ICRT, whose definition is reviewed in Section 3.2.1, is the asymptotic cone of the BHP_θ around its root, implying $\mathsf{BHP}_\theta \to \mathsf{ICRT}$ in law as $\theta \to \infty$. In particular, formally, we may think of the BHP_∞ as the ICRT. In view of Theorem 2, it is therefore natural to expect that the ICRT appears also as the scaling limit of the UIHPQ_{p_n}, provided θ in the definition of p_n is replaced by a sequence $\theta_n \to \infty$, that is, if $a_n^2(1 - 2p_n) \to \infty$ as $n \to \infty$. This is indeed the case.

Theorem 3. *Let $(a_n, n \in \mathbb{N})$ be a sequence of positive reals with $a_n \to \infty$. Let $(p_n, n \in \mathbb{N}) \subset [0, 1/2]$ be a sequence satisfying*

$$a_n^2(1 - 2p_n) \to \infty \quad \text{as } n \to \infty.$$

Then, in the sense of the local Gromov-Hausdorff topology as $n \to \infty$,

$$a_n^{-1} \cdot \mathsf{UIHPQ}_{p_n} \xrightarrow{(d)} \mathsf{ICRT}.$$

As special cases of the previous two theorems, we mention

Corollary 1. *Let $p \in [0, 1/2]$, and let $(a_n, n \in \mathbb{N})$ be a sequence of positive reals with $a_n \to \infty$. Then, in the sense of the local Gromov-Hausdorff topology as $n \to \infty$,*

$$a_n^{-1} \cdot \mathsf{UIHPQ}_p \xrightarrow{(d)} \begin{cases} \mathsf{ICRT} & \text{if} \quad 0 \leqslant p < 1/2 \\ \mathsf{BHP} & \text{if} \quad p = 1/2 \end{cases}.$$

For the family $(\mathbb{H}_\alpha)_\alpha$ of half-planar triangulations studied in [5, 37], convergence towards the ICRT in the subcritical regime $\alpha < 2/3$ is conjectured in [37, Section 2.1.2].



Figure 2: Illustration of various convergences explaining the connections between the spaces UIHPQ_p, BHP_θ and $\mathsf{BD}_{T,\sigma}$. For simplicity, the cases $\theta = 0$ and $\theta = \infty$ are left out. The top-most horizontal convergence represents [12, Theorem 1] and holds for $T, \sigma > 0$ fixed. If the volume T of $\mathsf{BD}_{T,\sigma}$ is blown up and the perimeter σ grows linearly in T such that $\sigma(T) \sim \theta T$, the space BHP_θ appears as the distributional local Gromov-Hausdorff limit of the disks $\mathsf{BD}_{T,\sigma(T)}$ around their roots ([8, Corollary 3.17]). On the other hand, BHP_θ is approximated by uniform quadrangulations $Q_n^{\sigma_n}$ ([8, Theorem 3.4]), or by the UIHPQ_p when $p = p(a_n, \theta)$ depends in the right way on θ and a_n (Theorem 2). The UIHPQ_p for fixed $p \in [0, 1/2]$ in turn arises as the local limit of $Q_n^{\sigma_n}$, provided the boundary lengths are properly chosen (Theorem 1).

Remark 4. We stress that the spaces BHP_θ can also be understood as Gromov-Hausdorff scaling limits of uniform quadrangulations $Q_n^{\sigma_n} \in \mathcal{Q}_n^{\sigma_n}$; see [8, Theorems 3.3, 3.4, 3.5]. More specifically, the BHP_θ for $\theta \in (0, \infty)$ arises when $\sqrt{n} \ll \sigma_n \ll n$ and the graph metric is rescaled by a factor a_n^{-1} satisfying $3\sigma_n a_n^2/(4n) \to \theta$ as n tends to infinity. The Brownian half-plane BHP_0 corresponding to the choice $\theta = 0$ appears more generally when $1 \ll \sigma_n \ll n$ and $1 \ll a_n \ll \min\{\sqrt{\sigma_n}, \sqrt{n/\sigma_n}\}$. Finally, the ICRT corresponding to $\theta = \infty$ appears when $\sigma_n \gg \sqrt{n}$ and $\max\{1, \sqrt{n/\sigma_n}\} \ll a_n \ll \sqrt{\sigma_n}$.

We may as well view the spaces BHP_θ as local scaling limits around the roots of the so-called Brownian disks $\mathsf{BD}_{T,\sigma}$ of volume $T > 0$ and perimeter $\sigma > 0$ introduced in [12]. More concretely, it was proved in [8, Corollaries 3.17, 3.18] that when both T and $\sigma = \sigma(T)$

tend to infinity such that $\sigma(T)/T \to \theta \in [0,\infty]$, then the BHP_θ is the local Gromov-Hausdorff limit in law of the disk $\mathsf{BD}_{T,\sigma(T)}$ around a boundary point chosen according to the boundary measure of the latter. Figure 2 depicts some convergences involving the families UIHPQ_p and BHP_θ.

2.3 Tree structure

We will prove that for $p < 1/2$, the UIHPQ_p can be represented as a collection of independent finite quadrangulations with a simple boundary glued along a tree structure. The tree structure is encoded by the looptree associated to a two-type version of Kesten's tree, and the finite quadrangulations are distributed according to the Boltzmann distribution $\widehat{\mathbb{P}}_g^\sigma$ on quadrangulations with a simple boundary of size 2σ. Precise definitions of the encoding objects are postponed to Section 3.

For $0 \leqslant p \leqslant 1/2$, let $g_p = p(1-p)/3$ and $z_p = (1-p)/4$. Let $F(g,z)$ be the partition function of the Boltzmann measure on finite rooted quadrangulations with a boundary, with weight g per inner face and $\sqrt{z}$ per boundary edge. Let moreover $\widehat{F}_k(g)$ be the partition function of the Boltzmann measure on finite rooted quadrangulations with a simple boundary of perimeter $2k$, with weight g per inner face.

We introduce two probability measures $\mu_\circ$ and $\mu_\bullet$ on $\mathbb{N}_0$ by setting

$$\mu_\circ(k) = \frac{1}{F(g_p,z_p)}\left(1-\frac{1}{F(g_p,z_p)}\right)^k, \quad k \in \mathbb{N}_0,$$
$$\mu_\bullet(2k+1) = \frac{1}{F(g_p,z_p)-1}\left[z_p F^2(g_p,z_p)\right]^{k+1}\widehat{F}_{k+1}(g_p), \quad k \in \mathbb{N}_0,$$

with $\mu_\bullet(k) = 0$ if k even. Exact expressions for $F(g_p,z_p)$ and $\widehat{F}_{k+1}(g_p)$ are given in (18) and (19) below. The fact that $\mu_\bullet$ is a probability distribution is a consequence of Identity (2.8) in [15]. We will prove in Lemma 11 that the pair $(\mu_\circ,\mu_\bullet)$ is critical for $0 \leqslant p < 1/2$, in the sense that the product of their respective means equals one, and subcritical if $p = 1/2$, meaning that the product of their means is strictly less than one. Moreover, both measures have small exponential moments. Our main result characterizing the structure of the UIHPQ_p for $0 \leqslant p < 1/2$ is the following.

Theorem 4. *Let $0 \leqslant p < 1/2$, and let $\mathrm{Loop}(\mathcal{T}_\infty)$ be the infinite looptree associated to Kesten's two-type tree $\mathcal{T}_\infty(\mu_\circ,\mu_\bullet)$. Glue into each inner face of $\mathrm{Loop}(\mathcal{T}_\infty)$ of degree 2σ an independent Boltzmann quadrangulations with a simple boundary distributed according to $\widehat{\mathbb{P}}_{g_p}^\sigma$. Then, the resulting infinite quadrangulation is distributed as the UIHPQ_p.*

The gluing operation fills in each (rooted) loop a finite-size quadrangulation with a simple boundary, which has the same perimeter as the loop. The two boundaries are glued together, such that the root edges of the loop and the quadrangulation get identified; see Remark 8. Figure 3 depicts the above representation of the UIHPQ_p in the case $0 < p < 1/2$, as well as the borderline cases $p = 0$ and $p = 1/2$. The branching structure of the standard $\mathsf{UIHPQ} = \mathsf{UIHPQ}_{1/2}$ has been investigated by Curien and Miermont [25]. They show that the UIHPQ can be seen as the uniform infinite half-planar quadrangulation with a simple boundary (represented by the big white semicircle in Figure 3), together with a collection



Figure 3: Schematic representation of the UIHPQ_p for $p \in [0, 1/2]$. On the left: The UIHPQ_0, that is, Kesten's tree associated to the critical geometric offspring distribution $\mu_{1/2}$. On the right: The standard uniform infinite half-planar quadrangulation UIHPQ with a general boundary. The white parts are understood to be filled in with quadrangulations, the big white semicircle representing the half-plane. In the middle: The UIHPQ_p with skewness parameter p. The white parts represent the (finite-size) quadrangulations with a simple boundary which are glued into the loops of the infinite looptree $\mathrm{Loop}(\mathcal{T}_\infty)$ associated to a two-type version $\mathcal{T}_\infty(\mu_\circ, \mu_\bullet)$ of Kesten's tree.

of finite-size quadrangulations with a general boundary, which are attached to the infinite simple boundary.

Remark 5. In the case $p = 0$, the above theorem can be seen as a restatement of Proposition 1. Indeed, in this case, one finds that $\mu_\circ = \mu_{1/2}$ is the critical geometric probability law, and $\mu_\bullet$ is the Dirac-distribution δ_1. By construction, all the inner faces of $\mathrm{Loop}(\mathcal{T}_\infty)$ have then degree 2, and the gluing of a Boltzmann quadrangulation distributed according to $\widehat{\mathbb{P}}^1_{g_0=0}$ simply amounts to close the face, by identifying its edges. One finally recovers Kesten's (one-type) tree associated to the offspring law $\mu_{1/2}$, as already found in Proposition 1.

Remark 6. In [9], it has been proved that geodesics in the standard UIHPQ intersect both the left and right part of the boundary infinitely many times (see [9, Section 2.3.3] for the exact terminology). However, up to removing finite quadrangulations that hang off from the boundary, the UIHPQ has the topology of a half-plane. Consequently, left and right parts of the boundary intersect only finitely many times. The branching structure described in Theorem 4 implies that the left and right parts of the boundary of the UIHPQ_p for $p < 1/2$ have *infinitely* many intersection points. As a consequence, any infinite self-avoiding path intersects both boundaries infinitely many times.

Our tree-like description of the UIHPQ_p for $0 \leqslant p < 1/2$ readily implies that simple random walk on the UIHPQ_p is recurrent. For $p = 0$, this result is due to Kesten [30].

Corollary 2. *Let $0 \leqslant p < 1/2$. Almost surely, simple random walk on the* UIHPQ_p *is recurrent.*

Somewhat informally, the tree structure describing the UIHPQ_p in the case $p < 1/2$ shows that there is an essentially unique way for the random walk to move to infinity. Said otherwise, the walk reduces essentially to a random walk on the half-line reflected at the origin, which is, of course, recurrent. We give a precise proof in terms of electric networks in Section 6.

Remark 7. As far as the standard uniform infinite half-planar quadrangulation UIHPQ corresponding to $p = 1/2$ is concerned, Angel and Ray [6] prove recurrence of the triangular analog with a simple boundary, the half-plane UIPT. They construct a full-plane extension of the half-plane UIPT using a decomposition into layers and then adapt the methods of Gurel-Gurevich and Nachmias [26], and Benjamini and Schramm [10]. It is believed that the arguments of [6] can be extended to the UIHPQ, too. Ray proves in [37] of recurrence of the half-plane models $\mathbb{H}_\alpha$ when $\alpha < 2/3$. In [13], Björnberg and Stefánsson prove that the (local) limit of bipartite Boltzmann planar maps is recurrent, for every choice of the weight sequence.

We believe that the mean displacement of a random walker after n steps on the UIHPQ_p for $p < 1/2$ is of order $n^{1/3}$, as for Kesten's tree (case $p = 0$). We will not pursue this further in this paper.

Let us finally mention another consequence of Theorem 4 concerning percolation thresholds. See, e.g., [4] for the terminology of Bernoulli percolation on random lattices.

Corollary 3. *Let $0 \leqslant p < 1/2$. The critical thresholds for Bernoulli site, bond and face percolation on the* UIHPQ_p *are almost surely equal to one.*

Therefore, percolation on the UIHPQ_p changes drastically depending on whether the skewness parameter p (not to be confused the the percolation parameter) is less or equal to $1/2$: In the standard $\mathsf{UIHPQ} = \mathsf{UIHPQ}_{1/2}$, the critical thresholds are known to be $5/9$ for site percolation, see [38], and $1/3$ for edge percolation and $3/4$ for face percolation, see [4]. The proof of the corollary follows immediately from Theorem 4.

3 Random half-planes and trees

In this section, we begin with a review of the one-parameter family of Brownian half-planes BHP_θ, $\theta \geqslant 0$, introduced in [8] (see also [27] for the case $\theta = 0$).

We then gather certain concepts around trees, which play an important role throughout this paper. We properly define the ICRT, two-type Galton-Watson trees and Kesten's infinite versions thereof, looptrees and the so-called tree of components.

3.1 The Brownian half-planes BHP_θ

We need some preliminary notation. Given a function $f = (f_t, t \in \mathbb{R})$, we set $\underline{f}_t = \inf_{[0,t]} f$ for $t \geqslant 0$ and $\underline{f}_t = \inf_{(-\infty,t]} f$ for $t < 0$. Moreover, if $f = (f_t, t \geqslant 0)$ is a real-valued function

indexed by the positive reals, its Pitman transform $\pi(f)$ is defined by

$$\pi(f)_t = f_t - 2\underline{f}_t.$$

In case $B = (B_t, t \geqslant 0)$ is a standard one-dimensional Brownian motion, its Pitman transform $\pi(B) = (\pi(B)_t, t \geqslant 0)$ is equal in law to a three-dimensional Bessel process, which has in turn the law of the modulus of a three-dimensional Brownian motion.

Now fix $\theta \in [0, \infty)$. The Brownian half-plane BHP_θ with skewness θ is defined in terms of its contour and label processes $X^\theta = (X^\theta_t, t \in \mathbb{R})$ and $W^\theta = (W^\theta_t, t \in \mathbb{R})$. They are characterized as follows.

- $(X^\theta_t, t \geqslant 0)$ has the law of a one-dimensional Brownian motion $B = (B_t, t \geqslant 0)$ with drift $-\theta$ and $B_0 = 0$, and $(X^\theta_{-t}, t \geqslant 0)$ has the law of the Pitman transform of an independent copy of B.
- Given X^θ, the (label) function W^θ has same distribution as $(\gamma_{-\underline{X}^\theta_t} + Z^\theta_t, t \in \mathbb{R})$, where
  - given X^θ, $Z^\theta = (Z^\theta_t, t \in \mathbb{R}) = Z^{X^\theta - \underline{X}^\theta}$ is a continuous modification of the centered Gaussian process with covariances given by

$$\mathbb{E}\left[Z^\theta_s Z^\theta_t\right] = \min_{[s\wedge t, s\vee t]} X^\theta - \underline{X}^\theta,$$

  - $(\gamma_x, x \in \mathbb{R})$ is a two-sided Brownian motion with $\gamma_0 = 0$ and scaled by the factor $\sqrt{3}$, independent of Z^θ.

The process Z^θ is usually called the (head of the) random snake driven by $X^\theta - \underline{X}^\theta$, see [32] for more on this. Next, we define two pseudo-metrics d_{X^θ} and d_{W^θ} on $\mathbb{R}$,

$$d_{X^\theta}(s,t) = X^\theta_s + X^\theta_t - 2\min_{[s\wedge t, s\vee t]} X^\theta, \quad \text{and} \quad d_{W^\theta}(s,t) = W^\theta_s + W^\theta_t - 2\min_{[s\wedge t, s\vee t]} W^\theta.$$

The pseudo-metric D_θ associated to BHP_θ is defined as the maximal pseudo-metric d on $\mathbb{R}$ satisfying $d \leqslant d_{W^\theta}$ and $\{d_{X^\theta} = 0\} \subseteq \{D_\theta = 0\}$. According to Chapter 3 of [18], it admits the expression $(s, t \in \mathbb{R})$

$$D_\theta(s,t) = \inf\left\{ \sum_{i=1}^k d_{W^\theta}(s_i, t_i) : \begin{array}{l} k \in \mathbb{N},\ s_1, \ldots, s_k, t_1, \ldots, t_k \in \mathbb{R}, s_1 = s, t_k = t, \\ d_{X^\theta}(t_i, s_{i+1}) = 0 \text{ for every } i \in \{1, \ldots, k-1\} \end{array} \right\}.$$

Definition 1. The Brownian half-plane BHP_θ has the law of the pointed metric space $(\mathbb{R}/\{D_\theta = 0\}, D_\theta, \rho_\theta)$, with the distinguished point ρ_θ is given by the equivalence class of 0.

Note that D_θ stands here also for the induced metric on the quotient space. It follows from standard scaling properties of X^θ and W^θ that for $\lambda > 0$, $\lambda \cdot \mathsf{BHP}_\theta =_d \mathsf{BHP}_{\theta/\lambda^2}$. In particular, BHP_0 is scale-invariant. It was shown in [8] that for every $\theta \geqslant 0$, BHP_θ has a.s. the topology of the closed half-plane $\overline{\mathbb{H}} = \mathbb{R} \times \mathbb{R}_+$.

3.2 Random trees and some of their properties

3.2.1 The infinite continuum random tree ICRT

Introduced by Aldous in [2], the ICRT is a random rooted real tree that forms the non-compact analog of the usual continuum random tree CRT. Consider the stochastic process $(X_t, t \in \mathbb{R})$ such that $(X_t, t \geqslant 0)$ and $(X_{-t}, t \geqslant 0)$ are two independent one-dimensional standard Brownian motions started at zero. Define on $\mathbb{R}$ the pseudo-metric

$$d_X(s,t) = X_s + X_t - 2 \min_{[s\wedge t, s\vee t]} X.$$

Definition 2. The ICRT is the continuum random real tree $\mathcal{T}_X$ coded by X, i.e., the ICRT has the law of the pointed metric space $(\mathcal{T}_X, d_X, [0])$, where $\mathcal{T}_X = \mathbb{R}/\{d_X = 0\}$, and the distinguished point is given by the equivalence class of 0.

The ICRT is scale-invariant, meaning that $\lambda \cdot \mathsf{ICRT} =_d \mathsf{ICRT}$ for $\lambda > 0$, and invariant under re-rooting. We remark that the ICRT is often defined in terms of two independent three-dimensional Bessel processes $(X_t, t \geqslant 0)$ and $(X_{-t}, t \geqslant 0)$. Since the Pitman transform π turns a Brownian motion into a three-dimension Bessel processes, it is readily seen that both definitions give rise to the same random tree.

3.2.2 (Sub)critical (two-type) Galton-Watson trees

We recall the formalism of (finite or infinite) plane trees, i.e., rooted ordered trees. The size $|\mathsf{t}| \in \mathbb{N}_0 \cup \{\infty\}$ of t is given by its number of edges, and we shall write $\mathbb{T}_f$ for the set of all finite plane trees.

We will often use the fact that if GW_ν denotes the law of a Galton-Watson tree with critical or subcritical offspring distribution ν, then

$$\mathsf{GW}_\nu(\mathsf{t}) = \prod_{u \in V(\mathsf{t})} \nu(k_u(\mathsf{t})), \quad t \in \mathbb{T}_f, \tag{1}$$

where for $u \in V(\mathsf{t})$, $k_u(\mathsf{t})$ is the number of offspring of vertex u. See, for example, [33, Proposition 1.4]). In the case where $\nu = \mu_p$ is the geometric offspring distribution of parameter $p \in [0, 1/2]$, (1) becomes

$$\mathsf{GW}_{\mu_p}(\mathsf{t}) = p^{|\mathsf{t}|}(1-p)^{|\mathsf{t}|+1}. \tag{2}$$

From the last display, the connection to random walks is apparent. Namely, let $(S^{(p)}(m), m \in \mathbb{N}_0)$ be a random walk on the integers starting from $S^{(p)}(0) = 0$ with increments distributed according to $p\delta_1 + (1-p)\delta_{-1}$. Define the first hitting time of -1,

$$T_{-1}^{(p)} = \inf\{m \in \mathbb{N} : S^{(p)}(m) = -1\}.$$

Then it is readily deduced from (2) that the size $|\mathsf{t}|$ of t under GW_{μ_p} and $(T_{-1}^{(p)} - 1)/2$ are equal in distribution; see, e.g., [36, Section 6.3].

Given a finite or infinite plane tree, it will be convenient to say that vertices at even height of t are white, and those at odd height are black. We use the notation $\mathsf{t}_\circ$ and $\mathsf{t}_\bullet$ for the associated subset of vertices. We next define two-type Galton-Watson trees associated to a pair $(\nu_\circ, \nu_\bullet)$ of probability measures on $\mathbb{N}_0$.

Definition 3. The *two-type Galton-Watson tree* with a pair of offspring distributions $(\nu_\circ, \nu_\bullet)$ is the random plane tree such that vertices at even height have offspring distribution $\nu_\circ$, vertices at odd height have offspring distribution $\nu_\bullet$, and the numbers of children of the different vertices are independent.

More formally, if $\mathsf{GW}_{\nu_\circ,\nu_\bullet}$ denotes the law of such a tree, then

$$\mathsf{GW}_{\nu_\circ,\nu_\bullet}(\mathsf{t}) = \prod_{u\in\mathsf{t}_\circ} \nu_\circ(k_u(\mathsf{t})) \prod_{u\in\mathsf{t}_\bullet} \nu_\bullet(k_u(\mathsf{t})), \quad t \in \mathbb{T}_f.$$

In this context, the pair $(\nu_\circ, \nu_\bullet)$ is said to be *critical* if and only if the mean vector $(m_\circ, m_\bullet)$ satisfies $m_\circ m_\bullet = 1$.

3.2.3 Kesten's tree and its two-type version

We next briefly review critical Galton-Watson trees conditioned to survive; see [30] or [35], and [39] for the multi-type case.

Proposition 3 (Theorem 3.1 in [39])**.** *Let* GW *be the law of a critical (either one or two-type) Galton-Watson tree. For every* $n \in \mathbb{N}$*, assume that* $\mathsf{GW}(\{\#V(\mathsf{t}) = n\}) > 0$*, and let* $\mathcal{T}_n$ *be a tree with law* GW *conditioned to have* n *vertices. Then, we have the local convergence for the metric* d_{map} *as* $n \to \infty$ *to a random infinite tree* $\mathcal{T}_\infty$*,*

$$\mathcal{T}_n \xrightarrow{(d)} \mathcal{T}_\infty.$$

In the case $\mathsf{GW} = \mathsf{GW}_\nu$ for ν a critical one-type offspring distribution, $\mathcal{T}_\infty$ is often called *Kesten's tree* associated to ν, and simply Kesten's tree if $\nu = \mu_{1/2}$. We will use the same terminology if $(\nu_\circ, \nu_\bullet)$ is a critical pair of offspring distributions and $\mathsf{GW} = \mathsf{GW}_{\nu_\circ,\nu_\bullet}$. In this case, we write $\mathcal{T}_\infty(\nu_\circ, \nu_\bullet)$ for Kesten's tree associated to $(\nu_\circ, \nu_\bullet)$. Note that the condition $\mathsf{GW}(\{\#V(\mathsf{t}) = n\}) > 0$ can be relaxed, provided we can find a subsequence along which this condition is satisfied.

Galton-Watson trees conditioned to survive enjoy an explicit construction, which we briefly recall for the two-type case. Details can be found in [39]. Let $(\nu_\circ, \nu_\bullet)$ be a critical pair of offspring distributions with mean $(m_\circ, m_\bullet)$, and recall that the size-biased distributions $\bar{\nu}_\circ$ and $\bar{\nu}_\bullet$ are defined by

$$\bar{\nu}_\circ(k) = \frac{k\nu_\circ(k)}{m_\circ} \quad \text{and} \quad \bar{\nu}_\bullet(k) = \frac{k\nu_\bullet(k)}{m_\bullet}, \quad k \in \mathbb{N}_0.$$

Kesten's tree $\mathcal{T}_\infty$ associated to $(\nu_\circ, \nu_\bullet)$ is an infinite locally finite (two-type) tree that has a.s. a unique infinite self-avoiding path called the *spine*. It is constructed as follows. The root vertex (white) is the first vertex on the spine. It has size-biased offspring distribution $\bar{\nu}_\circ$. Among its offspring, a child (black) is chosen uniformly at random to be the second vertex on the spine. It has size-biased offspring distribution $\bar{\nu}_\bullet$, and a child (white) chosen uniformly at random among its offspring becomes the third vertex on the spine. The spine is constructed by iterating this procedure.

The construction of the tree is completed by specifying that vertices at even (resp. odd) height lying not on the spine have offspring distribution $\nu_\circ$ (resp. $\nu_\bullet$), and that the numbers of offspring of the different vertices are independent.

The construction is similar in the mono-type case. In the particular case when $\nu = \mu_{1/2}$ is the geometric distribution with parameter $1/2$, Kesten's tree can be represented by an infinite half-line (isomorphic to $\mathbb{N}$) and a collection of independent Galton-Watson trees with law $\mathsf{GW}_{\mu_{1/2}}$ grafted to the left and to the right of every vertex on the spine; see, for instance, [28, Example 10.1]. We will exploit this representation in our proof of Proposition 1.

3.2.4 Random looptrees

Our description of the UIHPQ_p in Theorem 4 makes use of so-called *looptrees*, which were introduced in [22]. A looptree can informally be seen as a collection of loops glued along a tree structure. The following presentation is inspired by [23, Section 2.3]. We use, however, slightly different definitions which are better suited to our purpose. In particular, given a plane tree t, we will only replace vertices $v \in V(\mathsf{t}_\bullet)$ at *odd* height by loops of length $\deg(u)$. Consequently, several loops may be attached to one and the same vertex (at even height).

Let us now make things more precise. Let t be a finite plane tree, and recall that vertices at even height are white, and those at odd height are black (with respective subsets of vertices $\mathsf{t}_\circ$ and $\mathsf{t}_\bullet$). We associate to t a rooted looptree $\mathrm{Loop}(\mathsf{t})$ as follows. Around every (black) vertex in $\mathsf{t}_\bullet$, we connect its incident white vertices in cyclic order, so that they form a loop. Then $\mathrm{Loop}(\mathsf{t})$ is the planar map obtained from erasing the black vertices and the edges of t. We root $\mathrm{Loop}(\mathsf{t})$ at the edge connecting the origin of t to the last child of its first sibling in t; see Figure 4.

The reverse application associates to a looptree l a plane tree, which we call the *tree of components* $\mathrm{Tree}(\mathsf{l})$. In order to obtain $\mathrm{Tree}(\mathsf{l})$ from l, we add a new vertex in every internal face of l and connect this vertex to all the vertices of the face. $\mathrm{Tree}(\mathsf{l})$ is rooted at the corner adjacent to the target of the root edge of l. The root edge of $\mathrm{Tree}(\mathsf{l})$ connects the origin of l to the new vertex added in the face incident to the left side of the root edge of l.



Figure 4: A looptree and the associated tree of components.

The procedures Tree and Loop extend to infinite but locally finite trees, by considering the consistent sequence of maps $\{B_{2k}(\mathsf{t}) : k \in \mathbb{N}_0\}$. We will be interested in the random infinite looptree associated to Kesten's two-type tree.

Definition 4. If $(\nu_\circ, \nu_\bullet)$ is a critical pair of offspring laws and $\mathcal{T}_\infty$ the corresponding Kesten's tree, we call the random infinite looptree $\mathrm{Loop}(\mathcal{T}_\infty)$ *Kesten's looptree* associated to $\mathcal{T}_\infty$.

Note that a formal way to construct $\mathrm{Loop}(\mathcal{T}_\infty)$ is to define it as the local limit of $\mathrm{Loop}(\mathcal{T}_n)$, where $\mathcal{T}_n$ is a two-type Galton-Watson tree with offspring distribution $(\nu_\circ, \nu_\bullet)$ conditioned to have n vertices.

Remark 8. In a looptree l, every loop is naturally rooted at the edge whose origin is the closest vertex to the origin of l, such that the outer face of l lies on the right of that edge. The gluing of a (rooted) quadrangulation with a simple boundary of perimeter 2σ into a loop of the same length is then determined by the convention that the root edge of the quadrangulation is glued on the root edge of the loop.

4 Construction of the UIHPQ_p

A Schaeffer-type bijection due to Bouttier, Di Francesco and Guitter [14] encodes quadrangulations with a boundary in terms of labeled trees that are attached to a bridge. We shall first describe a bijective encoding of finite-size planar quadrangulations, and then extend it to infinite quadrangulations with an infinite boundary. This will allow us to construct and define the UIHPQ_p for $p \in [0, 1/2]$ in terms of the encoding objects, which we define first.

4.1 The encoding objects

We briefly review well-labeled trees, forests, bridges and contour and label functions. Our notation bears similarities to [25, 19, 8], differs, however, at some places. Each of these references already contains the construction of the standard UIHPQ.

4.1.1 Forest and bridges

A *well-labeled tree* (t, ℓ) is a pair consisting of a finite rooted plane tree t and a labeling $(\ell(u))_{u\in V(\mathsf{t})}$ of its vertices $V(\mathsf{t})$ by integers, with the constraints that the root vertex receives label zero, and $|\ell(u) - \ell(v)| \leqslant 1$ if u and v are connected by an edge.

A *well-labeled forest* with $\sigma \in \mathbb{N}$ trees is a pair $(\mathfrak{f}, \mathfrak{l})$, where $\mathfrak{f} = (\mathsf{t}_0, \ldots, \mathsf{t}_{\sigma-1})$ is a sequence of σ rooted plane trees, and $\mathfrak{l} : V(\mathfrak{f}) \to \mathbb{Z}$ is a labeling of the vertices $V(\mathfrak{f}) = \cup_{i=0}^{\sigma-1} V(\mathsf{t}_i)$ such that for every $0 \leqslant i \leqslant \sigma - 1$, the pair $(\mathsf{t}_i, \mathfrak{l}{\restriction}V(\mathsf{t}_i))$ is a well-labeled tree. Similarly, a *well-labeled infinite forest* is a pair $(\mathfrak{f}, \mathfrak{l})$, where $\mathfrak{f} = (\mathsf{t}_i, i \in \mathbb{Z})$ is an infinite collection of rooted plane trees, together with a labeling $\mathfrak{l} : \cup_{i\in\mathbb{Z}} V(\mathsf{t}_i) \to \mathbb{Z}$ such that for each $i \in \mathbb{Z}$, the restriction of $\mathfrak{l}$ to $V(\mathsf{t}_i)$ turns t_i into a well-labeled tree.

A *bridge of length* 2σ for $\sigma \in \mathbb{N}$ is a sequence $\mathsf{b} = (\mathsf{b}(0), \mathsf{b}(1), \ldots, \mathsf{b}(2\sigma-1))$ of 2σ integers with $\mathsf{b}(0) = 0$ and $|\mathsf{b}(i+1) - \mathsf{b}(i)| = 1$ for $0 \leqslant i \leqslant 2\sigma - 1$, where we agree that $\mathsf{b}(2\sigma) = 0$. In a similar manner, an *infinite bridge* is a two-sided sequence $\mathsf{b} = (\mathsf{b}(i) : i \in \mathbb{Z})$ with $\mathsf{b}(0) = 0$ and $|\mathsf{b}(i+1) - \mathsf{b}(i)| = 1$ for all $i \in \mathbb{Z}$.

Given a bridge b, an index i for which $\mathsf{b}(i+1) = \mathsf{b}(i) - 1$ is called a *down-step* of b. The set of all down-steps of b is denoted $\mathsf{DS}(\mathsf{b})$. If b is a bridge of length 2σ, $\mathsf{DS}(\mathsf{b})$ has σ elements, and we write $\mathrm{d}^\downarrow_\mathsf{b}(i)$ for the ith largest element in $\mathsf{DS}(\mathsf{b})$, for $i = 1, \ldots, \sigma$. If b is an infinite bridge and $i \in \mathbb{N}$, $\mathrm{d}^\downarrow_\mathsf{b}(i)$ denotes the ith largest element in $\mathsf{DS}(\mathsf{b}) \cap \mathbb{N}_0$, and $\mathrm{d}^\downarrow_\mathsf{b}(-i)$ denotes the ith largest element in $\mathsf{DS}(\mathsf{b}) \cap \mathbb{Z}_{<0}$. If there is no danger of confusion, we write simply $\mathrm{d}^\downarrow$ instead of $\mathrm{d}^\downarrow_\mathsf{b}$.

The size of a forest $\mathfrak{f}$ is the number $|\mathfrak{f}| \in \mathbb{N}_0 \cup \{\infty\}$ of tree edges. If $\mathfrak{f} = (\mathsf{t}_0, \dots, \mathsf{t}_{\sigma-1})$ and $u \in V(\mathsf{t}_i)$, we write $\mathrm{H}_{\mathfrak{f}}(u)$ for the height of u in the tree t_i, i.e., the graph distance to the root of t_i. Moreover, $\mathcal{I}_{\mathfrak{f}}(u) = i$ denotes the index of the tree the vertex u belongs to. Both $\mathrm{H}_{\mathfrak{f}}$ and $\mathcal{I}_{\mathfrak{f}}$ extend in the obvious way to infinite forests. If it is clear which forest we are referring to, we drop the subscript $\mathfrak{f}$ in H and $\mathcal{I}$.

We let $\mathfrak{F}_\sigma^n = \{(\mathfrak{f}, \mathfrak{l}) : \mathfrak{f} \text{ has } \sigma \text{ trees and size } |\mathfrak{f}| = n\}$ be the set of all well-labeled forests of size n with σ trees and write $\mathfrak{F}_\infty$ for the set of all well-labeled infinite forests. The set of all bridges of length 2σ is denoted $\mathfrak{B}_\sigma$. As far as infinite bridges are concerned, it will be sufficient to consider only those bridges b which satisfy $\inf_{i\in\mathbb{N}} \mathsf{b}(i) = -\infty$ and $\inf_{i\in\mathbb{N}} \mathsf{b}(-i) = -\infty$, and we denote the set of them by $\mathfrak{B}_\infty$.

4.1.2 Contour and label function

We first consider the case $((\mathfrak{f}, \mathfrak{l}), \mathsf{b}) \in \mathfrak{F}_\sigma^n \times \mathfrak{B}_\sigma$ for some $n, \sigma \in \mathbb{N}$. By a slight abuse of notation, we write $\mathfrak{f}(0), \dots, \mathfrak{f}(2n+\sigma-1)$ for the contour exploration of $\mathfrak{f}$, that is, the sequence of vertices (with multiplicity) which we obtain from walking around the trees $\mathsf{t}_0, \dots, \mathsf{t}_{\sigma-1}$ of $\mathfrak{f}$, one after the other in the contour order. See the left side of Figure 5. We define the *contour function* of $(\mathfrak{f}, \mathfrak{l})$ by

$$C_{\mathfrak{f}}(j) = \mathrm{H}(\mathfrak{f}(j)) - \mathcal{I}(\mathfrak{f}(j)), \quad 0 \leqslant j \leqslant 2n + \sigma - 1.$$

Note that $C_{\mathfrak{f}}(2n + \sigma - 1) = \sigma - 1$, since the last visited vertex by the contour exploration is the root of $\mathsf{t}_{\sigma-1}$. We extend $C_{\mathfrak{f}}$ to $[0, 2n + \sigma]$ by first letting $C_{\mathfrak{f}}(2n + \sigma) = -\sigma$, and then by linear interpolation between integers, so that $C_{\mathfrak{f}}$ becomes a continuous real-valued function on $[0, 2n + \sigma]$ starting at zero and ending at $-\sigma$.

The *label function* associated to $((\mathfrak{f}, \mathfrak{l}), \mathsf{b})$ is obtained from shifting the vertex label $\mathfrak{l}(\mathfrak{f}(j))$ by the value of the bridge b evaluated at its $(\mathcal{I}(\mathfrak{f}(j)) + 1)$th down-step. Formally,

$$\mathfrak{L}_{\mathfrak{f}}(j) = \mathfrak{l}(\mathfrak{f}(j)) + \mathsf{b}\left(\mathrm{d}^{\downarrow}\left(\mathcal{I}(\mathfrak{f}(j)) + 1\right)\right), \quad 0 \leqslant j \leqslant 2n + \sigma - 1.$$

We let $\mathfrak{L}_{\mathfrak{f}}(2n + \sigma) = 0$ and again linearly interpolate between integer values, so that $\mathfrak{L}_{\mathfrak{f}}$ becomes an element of $\mathcal{C}([0, 2n + \sigma], \mathbb{R})$. Contour and label functions are depicted on the right side of Figure 5.

In the case $((\mathfrak{f}, \mathfrak{l}), \mathsf{b}) \in \mathfrak{F}_\infty \times \mathfrak{B}_\infty$, we explore the trees of $\mathfrak{f}$ in the following way: First, $(\mathfrak{f}(0), \mathfrak{f}(1), \dots)$ is the sequence of vertices of the contour paths of the trees $\mathsf{t}_i, i \in \mathbb{N}_0$, in the left-to-right order, starting from the root of t_0. Then, we let $(\mathfrak{f}(-1), \mathfrak{f}(-2), \dots)$ be the sequence of vertices of the contour paths $\mathsf{t}_{-1}, \mathsf{t}_{-2}, \dots$, in the counterclockwise or right-to-left order, starting from the root of t_{-1}; see the left side of Figure 6. Contour and label functions $C_{\mathfrak{f}}$ and $\mathfrak{L}_{\mathfrak{f}}$ are defined similarly to the finite case, namely

$$\begin{aligned}
C_{\mathfrak{f}}(j) &= \mathrm{H}(\mathfrak{f}(j)) - \mathcal{I}(\mathfrak{f}(j)), \quad j \in \mathbb{Z}, \\
\mathfrak{L}_{\mathfrak{f}}(j) &= \mathfrak{l}(\mathfrak{f}(j)) + \mathsf{b}\left(\mathrm{d}^{\downarrow}\left(\mathcal{I}(\mathfrak{f}(j)) + 1\right)\right), \quad j \in \mathbb{Z}_{\geqslant 0}, \\
\mathfrak{L}_{\mathfrak{f}}(j) &= \mathfrak{l}(\mathfrak{f}(j)) + \mathsf{b}\left(\mathrm{d}^{\downarrow}\left(\mathcal{I}(\mathfrak{f}(j))\right)\right), \quad j \in \mathbb{Z}_{<0}.
\end{aligned}$$

Note that the asymmetry in the definition of $\mathfrak{L}_{\mathfrak{f}}$ stems from the numbering of the trees. By linear interpolation between integer values, we interpret $C_{\mathfrak{f}}$, $\mathfrak{L}_{\mathfrak{f}}$, and sometimes also $\mathfrak{l}$, as continuous functions (from $\mathbb{R}$ to $\mathbb{R}$).



Figure 5: Contour and label functions $C_\mathfrak{f}$ and $\mathfrak{L}_\mathfrak{f}$ of an element $((\mathfrak{f},\mathfrak{l}),\mathfrak{b}) \in \mathfrak{F}_4^7 \times \mathfrak{B}_4$. The left side depicts the contour exploration of $\mathfrak{f}$. The labels on the vertices are given by $\mathfrak{L}_\mathfrak{f}(j)$, $j = 0,\dots,18$. Note that the values of $\mathfrak{b}$ at its four down-steps are equal to the values of $\mathfrak{L}_\mathfrak{f}$ at the tree roots: In this example, we have $\mathfrak{b}(\mathrm{d}^\downarrow(1)) = 0$, $\mathfrak{b}(\mathrm{d}^\downarrow(2)) = -1$, and $\mathfrak{b}(\mathrm{d}^\downarrow(3)) = \mathfrak{b}(\mathrm{d}^\downarrow(4)) = 1$. The red dots on the right indicate the encoding of a new tree.

4.2 The Bouttier-Di Francesco-Guitter mapping

We denote the set of all rooted pointed quadrangulations with n inner faces and 2σ boundary edges by

$$\mathcal{Q}_n^{\sigma,\bullet} = \{(\mathfrak{q}, v^\bullet) : \mathfrak{q} \in \mathcal{Q}_n^\sigma, v^\bullet \in V(\mathfrak{q})\},$$

where $v^\bullet$ stands for the distinguished pointed vertex. In the following part, we briefly recall the definition of the bijection $\Phi_n : \mathfrak{F}_\sigma^n \times \mathfrak{B}_\sigma \to \mathcal{Q}_n^{\sigma,\bullet}$ introduced in [14].

4.2.1 The encoding of finite quadrangulations

We represent an element $((\mathfrak{f},\mathfrak{l}),\mathfrak{b}) \in \mathfrak{F}_\sigma^n \times \mathfrak{B}_\sigma$ in the plane as follows. Firstly, we view $\mathfrak{b}$ as a cycle of length 2σ: We start from a distinguished vertex labeled $\mathfrak{b}(0) = 0$ and label the remaining $2\sigma-1$ vertices in the counterclockwise order by the values $\mathfrak{b}(1), \mathfrak{b}(2), \dots, \mathfrak{b}(2\sigma-1)$. Then we graft the trees $(\mathfrak{t}_0,\dots,\mathfrak{t}_{\sigma-1})$ of $\mathfrak{f}$ to the σ down-steps $0 \leqslant i_0 < i_1 < \cdots < i_{\sigma-1} \leqslant 2\sigma-1$ of $\mathfrak{b}$, such that $\mathfrak{t}_j$ is grafted on the vertex corresponding to the value $\mathfrak{b}(i_j)$, in the interior of the cycle. We do it in such a way that different trees do not intersect. The vertices of $\mathfrak{t}_j$ are equipped with their labels shifted by $\mathfrak{b}(i_j)$. Figure 7 illustrates this procedure.

We now build a rooted and pointed quadrangulation $(\mathfrak{q}, v^\bullet)$ out of $((\mathfrak{f},\mathfrak{l}),\mathfrak{b})$. First, we put an extra vertex $v^\bullet$ in the interior of the cycle representing $\mathfrak{b}$. The set of vertices of $\mathfrak{q}$ is given by the tree vertices $V(\mathfrak{f}) \cup \{v^\bullet\}$. As for the edges of $\mathfrak{q}$, we define for $0 \leqslant i \leqslant 2n+\sigma-1$ the successor $\mathrm{succ}(i) \in [0, 2n+\sigma-1] \cup \{\infty\}$ of i to be the first element k in the list $(i+1,\dots,2n+\sigma-1,0,\dots,i-1)$ (from left to right) which has label $\mathfrak{L}_\mathfrak{f}(k) = \mathfrak{L}_\mathfrak{f}(i) - 1$. If there is no such element, we put $\mathrm{succ}(i) = \infty$. We extend the contour exploration $\mathfrak{f}(0),\dots,\mathfrak{f}(2n+\sigma-1)$ of $\mathfrak{f}$ by setting $\mathfrak{f}(\infty) = v^\bullet$. We follow the exploration starting from the vertex $\mathfrak{f}(0)$ (which is the root of $\mathfrak{t}_0$) and draw for each $0 \leqslant i \leqslant 2n+\sigma-1$ an arc between $\mathfrak{f}(i)$ and $\mathfrak{f}(\mathrm{succ}(i))$, such that arcs do not cross. Except for the leaves, a vertex of $\mathfrak{f}$ is visited at least twice in the contour exploration, so that there are in general several arcs connecting the vertices $\mathfrak{f}(i)$ and $\mathfrak{f}(\mathrm{succ}(i))$. The edges of $\mathfrak{q}$ are given by all these arcs between the vertices $V(\mathfrak{f}) \cup \{v^\bullet\}$.



Figure 6: Contour and label functions $C_{\mathfrak{f}}$ and $\mathfrak{L}_{\mathfrak{f}}$ of an element $((\mathfrak{f},\mathfrak{l}),\mathfrak{b}) \in \mathfrak{F}_\infty \times \mathfrak{B}_\infty$. The left side depicts the two-sided contour exploration of $\mathfrak{f}$. The labels are given by $\mathfrak{L}_{\mathfrak{f}}(j)$, where now $j \in \mathbb{Z}$. The values of the infinite bridge $\mathfrak{b}$ at its first three down-steps to the right of 0 read here $\mathfrak{b}(\mathrm{d}^{\downarrow}(1)) = 2$, $\mathfrak{b}(\mathrm{d}^{\downarrow}(2)) = 1$ and $\mathfrak{b}(\mathrm{d}^{\downarrow}(3)) = 3$, while the first down-step to the left of zero has value $\mathfrak{b}(\mathrm{d}^{\downarrow}(-1)) = 0$. The arrows below the contour function indicate the direction of the encoding, and the red dots mark again the encoding of a new tree.

It only remains to root the quadrangulation. To that aim, we observe from Figure 7 that the 2σ boundary edges of $\mathfrak{q}$ are in a order-preserving correspondence with the 2σ cycle edges. We root $\mathfrak{q}$ at the edge corresponding to the first edge of the cycle (starting from the distinguished edge, in the clockwise order), oriented in such a way that the face of degree 2σ becomes the outer face (i.e., lies to the right of the root edge). Upon erasing the tree and cycle edges of the representation of $((\mathfrak{f},\mathfrak{l}),\mathfrak{b})$, and the vertices of $\mathfrak{b}$ corresponding to up-steps, we obtain a rooted pointed quadrangulation $(\mathfrak{q}, v^\bullet)$. A description of the reverse mapping $\Phi_n^{-1} : \mathcal{Q}_n^{\sigma,\bullet} \to \mathfrak{F}_\sigma^n \times \mathfrak{B}_\sigma$ can be found in [14] or [11].

4.2.2 The encoding of infinite quadrangulations

Recall that $\mathcal{Q}$ is the completion of the set of finite rooted quadrangulations with a boundary with respect to d_{map}. The aim of this section is to extend Φ_n to a mapping

$$\Phi : (\cup_{n,\sigma\in\mathbb{N}} \mathfrak{F}_\sigma^n \times \mathfrak{B}_\sigma) \cup (\mathfrak{F}_\infty \times \mathfrak{B}_\infty) \longrightarrow \mathcal{Q}.$$

We proceed as follows. If $((\mathfrak{f},\mathfrak{l}),\mathfrak{b}) \in \mathfrak{F}_\sigma^n \times \mathfrak{B}_\sigma$, we put $\Phi((\mathfrak{f},\mathfrak{l}),\mathfrak{b}) = \Phi_n((\mathfrak{f},\mathfrak{l}),\mathfrak{b})$. (We forget the distinguished vertex of $\Phi_n((\mathfrak{f},\mathfrak{l}),\mathfrak{b})$ and view the quadrangulation as an element in $\mathcal{Q}_n^\sigma \subset \mathcal{Q}$.)

Now assume $((\mathfrak{f},\mathfrak{l}),\mathfrak{b}) \in \mathfrak{F}_\infty \times \mathfrak{B}_\infty$. We consider the following representation of $((\mathfrak{f},\mathfrak{l}),\mathfrak{b})$ in the upper half-plane: First, we identify $\mathfrak{b}$ with the bi-infinite line obtained from connecting $i \in \mathbb{Z}$ to $i+1$ by an edge. Vertex i is labeled $\mathfrak{b}(i)$. We attach the trees $\mathfrak{t}(0), \mathfrak{t}(1), \ldots$ of $\mathfrak{f}$ to the down-steps of $\mathfrak{b}$ to the right of 0, and the trees $\mathfrak{t}(-1), \mathfrak{t}(-2), \ldots$ to the down-steps of $\mathfrak{b}$ to the left of -1, everything in the upper half-plane. Again, the labels in a tree are shifted by the underlying bridge label.

Similarly to the finite case, the vertex set of $\mathfrak{q} = \Phi((\mathfrak{f},\mathfrak{l}),\mathfrak{b})$ is given by $V(\mathfrak{f})$; here, we add no additional vertex. For specifying the edges, we let the successor $\mathrm{succ}_\infty(i)$ of $i \in \mathbb{Z}$ be the smallest number $k > i$ such that $\mathfrak{L}_{\mathfrak{f}}(k) = \mathfrak{L}_{\mathfrak{f}}(i) - 1$. Since by assumption $\inf_{i\in\mathbb{N}} \mathfrak{b}(i) = -\infty$, $\mathrm{succ}_\infty(i)$ is a finite number. We next connect the vertices $\mathfrak{f}(i)$ and $\mathfrak{f}(\mathrm{succ}_\infty(i))$ by an arc for any $i \in \mathbb{Z}$, such that the resulting map is planar. The arcs form the edges of the infinite



Figure 7: A representation of an element $((\mathfrak{f},\mathfrak{l}),\mathsf{b}) \in \mathfrak{F}_6^6 \times \mathfrak{B}_6$ in the plane and the associated rooted pointed quadrangulation $(\mathsf{q}, v^\bullet) = \Phi_n((\mathfrak{f},\mathfrak{l}),\mathsf{b})$. The distinguished vertex of the cycle is the down-most vertex labeled 0. The trees are grafted to the 6 down-steps of b (here, $\mathrm{d}^\downarrow(1) = 0$, $\mathrm{d}^\downarrow(2) = 1$, $\mathrm{d}^\downarrow(3) = 8$, $\mathrm{d}^\downarrow(4) = 10$, $\mathrm{d}^\downarrow(5) = 11$, and $\mathrm{d}^\downarrow(6) = 12$). The tree edges are indicated by the dashed lines in the interior of the cycle. Note that three trees (those above the first, fourth and sixth down-step) consist of a single vertex. The labels in a tree are shifted by the bridge value of the down-step above which the tree is attached. Note that the 12 boundary edges of the cycle are in a order-preserving correspondence with the 12 boundary edges of q. (The two edges of q which lie entirely in the outer face are counted twice.)

rooted quadrangulation q we are about to construct. In order to root the map, we observe that the bi-infinite line $\mathbb{Z}$ is in correspondence with the boundary edges of q, and we choose the edge corresponding to $\{0,1\}$ as the root edge of q (oriented such that the outer face lies to its right). A representation of $((\mathfrak{f},\mathfrak{l}),\mathsf{b})$ and of the resulting quadrangulation $\Phi((\mathfrak{f},\mathfrak{l}),\mathsf{b})$ is depicted in Figure 8.

4.3 Definition of the UIHPQ_p

We are now in position to construct the UIHPQ_p by means of the above mapping Φ applied to a (random) element in $\mathfrak{F}_\infty \times \mathfrak{B}_\infty$, which we introduce first.

Let t be a finite random plane tree. Conditionally on t, we assign to t a random uniform labeling ℓ of its vertices, so that the pair (t,ℓ) becomes a well-labeled tree. Namely, given t, we first equip each edge of t with an independent random variable uniformly distributed in $\{-1,0,1\}$. Then we define the label $\ell(u)$ of a vertex $u \in V(\mathsf{t})$ to be the sum over all labels along the unique (non-backtracking) path from the tree root to u.

We consider Galton-Watson trees with a (sub-)critical geometric offspring law μ_p of parameter p, $p \in [0,1/2]$, that is, $\mu_p(k) = p^k(1-p)$, $k \in \mathbb{N}_0$. If t is such a tree, we call it a *p-Galton-Watson tree*. Equipped with a random uniform labeling ℓ as described before, we say that the pair $(\mathsf{t}, (\ell(u))_{u\in V(\mathsf{t})})$ is a *uniformly labeled p-Galton-Watson tree*.

A *uniformly labeled infinite p-forest* is a random element $(\mathfrak{f}_\infty^{(p)}, \mathfrak{l}_\infty^{(p)})$ taking values in $\mathfrak{F}_\infty$, such that $(\mathsf{t}_i, \mathfrak{l}_\infty^{(p)} \restriction V(\mathsf{t}_i))$, $i \in \mathbb{Z}$, are independent uniformly labeled p-Galton-Watson trees.



Figure 8: The Bouttier-Di Francesco-Guitter mapping applied to an element $((\mathfrak{f}, \mathfrak{l}), \mathsf{b}) \in \mathfrak{F}_\infty \times \mathfrak{B}_\infty$. On the right hand side, the arcs connect the vertices $\mathfrak{f}(i)$ with $\mathfrak{f}(\mathrm{succ}_\infty(i))$, for $i \in \mathbb{Z}$. The other vertices and edges of the representation of $((\mathfrak{f}, \mathfrak{l}), \mathsf{b})$ on the left hand side do not appear in the quadrangulation. The oriented arc on the right indicated by an arrow represents the root edge of the map.

A *uniform infinite bridge* is a random element $\mathsf{b}_\infty = (\mathsf{b}_\infty(i), i \in \mathbb{Z})$ in $\mathfrak{B}_\infty$ such that b_∞ has the law of a two-sided simple symmetric random walk starting from $\mathsf{b}_\infty(0) = 0$. We stress that our wording differs from [8], where a uniform infinite bridge refers to a two-sided random walk with a geometric offspring law of parameter 1/2. See also Lemma 2 below.

Definition 5. Fix $p \in [0, 1/2]$. Let $(\mathfrak{f}_\infty^{(p)}, \mathfrak{l}_\infty^{(p)})$ be a uniformly labeled infinite p-forest, and independently of $(\mathfrak{f}_\infty^{(p)}, \mathfrak{l}_\infty^{(p)})$, let b_∞ be a uniform infinite bridge. Then the UIHPQ_p with skewness parameter p is given by the (rooted) random infinite quadrangulation $Q_\infty^\infty(p) = (V(Q_\infty^\infty(p)), d_{\mathrm{gr}}, \rho)$ with an infinite boundary, which is obtained from applying the Bouttier-Di Francesco-Guitter mapping Φ to $((\mathfrak{f}_\infty^{(p)}, \mathfrak{l}_\infty^{(p)}), \mathsf{b}_\infty)$. In case $p = 1/2$, we simply write Q_∞^∞, which denotes then the (standard) uniform infinite half-planar quadrangulation with a general boundary.

Remark 9. Let $\mathfrak{f}_\infty^{(p)}$ be the encoding forest of the UIHPQ_p. Instead of working with metric balls around the root vertex in the UIHPQ_p, it will – due to the specific construction of the latter – often be more practical to consider metric balls around the vertex corresponding to the tree root $\mathfrak{f}_\infty^{(p)}(0)$ in the UIHPQ_p. Similarly, if $Q_n^\sigma \in \mathcal{Q}_n^\sigma$ is a uniform quadrangulation and $\mathfrak{f}_n$ its encoding forest, it will be more natural to consider balls around $\mathfrak{f}_n(0)$ in Q_n^σ. Since the distance between $\mathfrak{f}_\infty^{(p)}(0)$ or $\mathfrak{f}_n(0)$ and the root of the map is stochastically bounded (it may also be zero), this makes no difference in terms of scaling limits whatsoever; see [8, Lemma 5.6]. We shall use the notation $B_r^{(0)}(Q_\infty^\infty(p))$ for the metric ball of radius r around $\mathfrak{f}_\infty^{(p)}(0)$ in the UIHPQ_p. Analogously, we define $B_r^{(0)}(Q_n^\sigma)$.

5 Proofs of the limit results

5.1 The UIHPQ_p as a local limit of uniform quadrangulations

In this part, we prove Theorem 1 and Proposition 1. We begin with the former. The case $p = 1/2$ has already been treated in [8], and the case $p = 0$ will be considered afterwards, so we first fix $0 < p < 1/2$ and let $(\sigma_n, n \in \mathbb{N})$ be a sequence of positive integers satisfying $\sigma_n = \frac{1-2p}{p} n + o(n)$. Recall that rooted pointed quadrangulations in $\mathcal{Q}^\bullet_{n,\sigma_n}$ are in one-to-one correspondence with elements in $\mathfrak{F}^n_{\sigma_n} \times \mathfrak{B}_{\sigma_n}$. For proving Theorem 1, the key step is to control the law of the first k trees in a forest $\mathfrak{f}_n$ chosen uniformly at random in $\mathfrak{F}^n_{\sigma_n}$, for k arbitrarily large but fixed. We will see in Lemma 1 below that their law is close to the law of k independent p-Galton-Watson trees when n is sufficiently large. Together with a convergence result of bridges (Lemma 2), this allows us to couple contour and label functions of $Q_n^{\sigma_n}$ and the UIHPQ_p, such that with high probability, we have equality of balls of a constant radius around the roots in $Q_n^{\sigma_n}$ and the UIHPQ_p, respectively. This readily implies the theorem.

We begin with the necessary control over the trees. Since the result on the tree convergence is of some interest on its own, we formulate an optimal version, which is stronger than we what need for mere local convergence as stated in Theorem 1. The exact formulation depends on the error term in the expression for σ_n. Let us put

$$\gamma_n = \max\left\{\sigma_n - \frac{1-2p}{p} n,\, 1\right\}. \tag{3}$$

Lemma 1. *Fix $0 < p < 1/2$, and let $(\sigma_n, n \in \mathbb{N})$ be a sequence of positive integers satisfying $\sigma_n = \frac{1-2p}{p} n + o(n)$. Define γ_n in terms of σ_n and p. Let $(\mathsf{t}_i)_{1 \leqslant i \leqslant \sigma_n}$ be a family of σ_n independent 1/2-Galton-Watson trees, and let $(\mathsf{t}_i^{(p)})_{1 \leqslant i \leqslant \sigma_n}$ be a family of σ_n independent p-Galton-Watson trees. Then, if $(k_n, n \in \mathbb{N})$ is a sequence of positive integers satisfying $k_n \leqslant \sigma_n$ and $k_n = o\left(\min\{n^{1/2}, n/\gamma_n\}\right)$ as $n \to \infty$, we have*

$$\lim_{n\to\infty} \left\| \mathrm{Law}\Big((\mathsf{t}_i)_{1\leqslant i\leqslant k_n} \,\Big|\, \sum_{i=1}^{\sigma_n} |\mathsf{t}_i| = n\Big) - \mathrm{Law}\left((\mathsf{t}_i^{(p)})_{1\leqslant i\leqslant k_n}\right) \right\|_{\mathrm{TV}} = 0.$$

Remark 10. We stress that if we only know $\gamma_n = o(n)$ as assumed in the statement of Theorem 1, we can at least choose k_n equals an (arbitrary) large constant $k \in \mathbb{N}$. This suffices in any case to show local convergence towards the UIHPQ_p; see Proposition 4 below. Lemma 1 may be seen as a complement to the results on coupling of trees in [8]; it treats a regime not considered in that work.

Proof. We write $\mathbb{P}_n$ for the conditional law of $(\mathsf{t}_i)_{1\leqslant i\leqslant k_n}$ given $\sum_{i=1}^{\sigma_n} |\mathsf{t}_i| = n$, and $\mathbb{Q}_n$ for the (unconditioned) law of $(\mathsf{t}_i^{(p)})_{1\leqslant i\leqslant k_n}$. Given a family $\mathfrak{f}$ of k_n trees, we write $\mathbf{v}(\mathfrak{f})$ for the sum of their sizes, i.e., the total number of edges. Note that

$$\mathrm{supp}(\mathbb{P}_n) = \mathrm{supp}(\mathbb{Q}_n) \cap \{\mathfrak{f} : \mathbf{v}(\mathfrak{f}) \leqslant n\}.$$

We now proceed in two steps. First, we show that for each $\varepsilon > 0$, there exists a constant $K > 0$ such that

$$\mathbb{Q}_n\left(\{\mathfrak{f} : \mathbf{v}(\mathfrak{f}) > Kk_n\}\right) \leqslant \varepsilon, \quad \mathbb{P}_n\left(\{\mathfrak{f} : \mathbf{v}(\mathfrak{f}) > Kk_n\}\right) \leqslant \varepsilon. \tag{4}$$

We then show that for large enough n, we have for any $\mathfrak{f} \in \mathrm{supp}(\mathbb{P}_n)$ of total size $\mathbf{v}(\mathfrak{f}) \leqslant Kk_n$,

$$1 - \varepsilon \leqslant \left| \frac{\mathbb{P}_n(\mathfrak{f})}{\mathbb{Q}_n(\mathfrak{f})} \right| \leqslant 1 + \varepsilon. \tag{5}$$

Clearly, (4) and (5) imply the claim of the lemma. We first prove (4). Let $(S^{(p)}(m), m \in \mathbb{N}_0)$ be a random walk on the integers starting from $S^{(p)}(0) = 0$ with increments distributed according to $p\delta_1 + (1-p)\delta_{-1}$. Set, for $\ell \in \mathbb{Z}$,

$$T_\ell^{(p)} = \inf\left\{m \in \mathbb{N} : S^{(p)}(m) = \ell\right\}.$$

Note that $S^{(p)}(m) + (1-2p)m$, $m \in \mathbb{N}_0$, is a martingale. We now use that the total size of k_n trees under $\mathbb{Q}_n$ and $(T^{(p)}_{-k_n} - k_n)/2$ are equal in distribution; see the discussion in Section 3.2.2. Applying Markov's inequality in the second and the optional stopping theorem in the third step, we obtain for K large enough

$$\mathbb{Q}_n\left(\mathbf{v} > Kk_n\right) = \mathbb{P}\left(T^{(p)}_{-k_n} > (2K+1)k_n\right) \leqslant \frac{\mathbb{E}[T^{(p)}_{-k_n}]}{(2K+1)k_n} = \frac{1}{(1-2p)(2K+1)} \leqslant \varepsilon.$$

For bounding the second probability in (4), we let $(S(m), m \in \mathbb{N}_0)$ be a simple symmetric random walk started from $S(0) = 0$ and write T_ℓ for its first hitting time of $\ell \in \mathbb{Z}$. Then

$$\mathbb{P}_n\left(\mathbf{v} > Kk_n\right) = \mathbb{P}\left(T_{-k_n} > Kk_n \,|\, T_{-\sigma_n} = 2n + \sigma_n\right).$$

Let us abbreviate $N = 2n + \sigma_n$, and $K_n = \lceil Kk_n \rceil$. We recall Kemperman's formula; see [36, Section 6.1]. Applying first the Markov property at time K_n and then Kemperman's formula to both the nominator and denominator gives, for large n,

$$\begin{aligned}
&\mathbb{P}\left(T_{-k_n} > K_n \,|\, T_{-\sigma_n} = N\right) = \mathbb{E}\left[\mathbf{1}_{\{T_{-k_n} > K_n\}} \frac{\mathbb{P}\left(T_{-\sigma_n} = N \,|\, S(K_n)\right)}{\mathbb{P}\left(T_{-\sigma_n} = N\right)}\right] \\
&= \mathbb{E}\left[\mathbf{1}_{\{T_{-k_n} > K_n\}} \frac{\sigma_n(\sigma_n + S(K_n))}{N(N-K_n)} \frac{\mathbb{P}\left(S(N) = -\sigma_n \,|\, S(K_n)\right)}{\mathbb{P}\left(S(N) = -\sigma_n\right)}\right] \\
&\leqslant 2\,\mathbb{P}\left(S(K_n) > -k_n \,|\, S(N) = -\sigma_n\right).
\end{aligned}$$

Let $(\tilde{S}(m), m \in \mathbb{N}_0)$ be the random walk starting from $\tilde{S}(0) = 0$ with steps

$$1 + \frac{\sigma_n}{N} \quad \text{with probability } \frac{1 - \sigma_n/N}{2}, \qquad -1 + \frac{\sigma_n}{N} \quad \text{with probability } \frac{1 + \sigma_n/N}{2}.$$

Clearly, $(\tilde{S}(m), m \in \mathbb{N}_0)$ is a martingale, and we find the relation

$$\mathbb{P}\left(S(K_n) \geqslant -k_n \,\middle|\, S(N) = -\sigma_n\right) = \mathbb{P}\left(\tilde{S}(K_n) \geqslant K_n \frac{\sigma_n}{N} - k_n \,\middle|\, \tilde{S}(N) = 0\right).$$

We now estimate

$$\begin{aligned}
&\mathbb{P}\left(\tilde{S}(K_n) \geqslant K_n \frac{\sigma_n}{N} - k_n \,\middle|\, \tilde{S}(N) = 0\right) \\
&\leqslant \mathbb{P}\left(\tilde{S}(N) \geqslant 0\right)^{-1} \mathbb{P}\left(\tilde{S}(K_n) \geqslant K_n \frac{\sigma_n}{N} - k_n\right) \leqslant 3 \frac{\mathbb{E}\left[\tilde{S}(K_n)^2\right]}{(K_n \frac{\sigma_n}{N} - k_n)^2} \leqslant \frac{12 K_n}{(K_n \frac{\sigma_n}{N} - k_n)^2}.
\end{aligned}$$

Here, in the next to last inequality, we have used Doob's inequality, as well as the bound $\mathbb{P}(\tilde{S}(N) \geqslant 0) \geqslant 1/3$ for large n, which is a direct consequence of the martingale central limit theorem. Since σ_n/N remains bounded away from zero (recall that $p < 1/2$), the last expression on the right hand side can be made arbitrarily small, upon choosing K large. This proves $\mathbb{P}_n(\mathbf{v} > Kk_n) \leqslant \varepsilon$ for K large enough, and hence (4) holds.

We turn to (5). First, observe that for a fixed $\mathfrak{f}$ in the support of $\mathbb{P}_n$, $\mathbb{P}_n(\mathfrak{f})$ is the probability to see k_n particular trees of total size $\mathbf{v}(\mathfrak{f})$ in a forest consisting of σ_n trees with total size n. An application of Kemperman's formula gives

$$\mathbb{P}_n(\mathfrak{f}) = \frac{\frac{\sigma_n - k_n}{2(n-\mathbf{v}(\mathfrak{f}))+\sigma_n-k_n} 2^{2(n-\mathbf{v}(\mathfrak{f}))+\sigma_n-k_n} \mathbb{P}\left(S(2(n-\mathbf{v}(\mathfrak{f}))+\sigma_n-k_n) = \sigma_n - k_n\right)}{\frac{\sigma_n}{2n+\sigma_n} 2^{2n+\sigma_n} \mathbb{P}\left(S(2n+\sigma_n) = \sigma_n\right)}.$$

Since $k_n \ll n$, we have for large $n \in \mathbb{N}$ and all families $\mathfrak{f}$ of k_n trees with $\mathbf{v}(\mathfrak{f}) \leqslant Kk_n$,

$$1 - \varepsilon \leqslant \left| \frac{\frac{\sigma_n - k_n}{2(n-\mathbf{v}(\mathfrak{f}))+\sigma_n-k_n} 2^{2(n-\mathbf{v}(\mathfrak{f}))+\sigma_n-k_n}}{\frac{\sigma_n}{2n+\sigma_n} 2^{2n+\sigma_n}} 2^{2\mathbf{v}(\mathfrak{f})+k_n} \right| \leqslant 1 + \varepsilon.$$

On the other hand, we know from (2) that $\mathbb{Q}_n(\mathfrak{f}) = p^{\mathbf{v}(\mathfrak{f})}(1-p)^{\mathbf{v}(\mathfrak{f})+k_n}$. Display (5) will therefore follow if we show that for large n and all $\mathfrak{f}$ with $\mathbf{v}(\mathfrak{f}) \leqslant Kk_n$,

$$1 - \varepsilon \leqslant \left| \frac{\mathbb{P}\left(S(2(n-\mathbf{v}(\mathfrak{f}))+\sigma_n-k_n) = \sigma_n - k_n\right)}{\mathbb{P}\left(S(2n+\sigma_n) = \sigma_n\right)} ((2p)^{-\mathbf{v}(\mathfrak{f})}(2(1-p))^{-(\mathbf{v}(\mathfrak{f})+k_n)} \right| \leqslant 1 + \varepsilon. \quad (6)$$

For a given $\mathfrak{f}$ with $\mathbf{v}(\mathfrak{f}) \leqslant Kk_n$, let us abbreviate $y_n = 2(n-\mathbf{v}(\mathfrak{f}))+\sigma_n-k_n$, $x_n = \sigma_n - k_n$, and $v_n = \mathbf{v}(\mathfrak{f})$. Clearly,

$$\frac{\mathbb{P}\left(S(2(n-v_n)+\sigma_n-k_n) = \sigma_n - k_n\right)}{\mathbb{P}\left(S(2n+\sigma_n) = \sigma_n\right)} = \frac{\binom{y_n}{\frac{y_n - x_n}{2}}}{\binom{2n+\sigma_n}{n}} 2^{2v_n+k_n}.$$

Combining the last two displays, it remains to show that

$$1 - \varepsilon \leqslant \left| \frac{y_n!(n+\sigma_n)!n!}{\frac{y_n-x_n}{2}!\frac{y_n+x_n}{2}!(2n+\sigma_n)!} p^{-v_n}(1-p)^{-(v_n+k_n)} \right| \leqslant 1 + \varepsilon.$$

The constants in the following error terms are uniform in the choice of $\mathfrak{f}$ satisfying $v_n = \mathbf{v}(\mathfrak{f}) \leqslant Kk_n$. By Stirling's formula and a rearrangement of the terms, we obtain

$$\frac{y_n!(n+\sigma_n)!n!}{\frac{y_n-x_n}{2}!\frac{y_n+x_n}{2}!(2n+\sigma_n)!} = (1+o(1)) \times \underbrace{\left(\frac{y_n}{2n+\sigma_n}\right)^{2n+\sigma_n} \left(\frac{n}{n-v_n}\right)^{n} \left(\frac{n+\sigma_n}{(y_n+x_n)/2}\right)^{n+\sigma_n}}_{=\mathrm{I}} \underbrace{\left(\frac{n-v_n}{y_n}\right)^{v_n} \left(\frac{(y_n+x_n)/2}{y_n}\right)^{v_n+k_n}}_{=\mathrm{II}}.$$

Recall that $k_n = o\left(\min\{n^{1/2}, n/\gamma_n\}\right)$ and $v_n \leqslant Kk_n$. We replace x_n and y_n by their values and obtain for the product I of the first three factors

$$\mathrm{I} = \exp\left(-(2v_n+k_n)\right) \exp\left((v_n+k_n)\right) \exp\left(v_n\right) \left(1 + O(k_n^2/n)\right) = 1 + o(1).$$

Recalling the particular form of σ_n for the product II of the last two factors, we arrive at

$$\mathrm{II} = p^{v_n}(1-p)^{v_n+k_n}\,(1+O(\gamma_n/n))^{2v_n+k_n} = p^{v_n}(1-p)^{v_n+k_n}\,(1+o(1))\,.$$

This proves (6) and hence the lemma. □

We continue with a convergence result for uniform bridges $\mathsf{b}_n \in \mathfrak{B}_{\sigma_n}$ towards b_∞.

Lemma 2. *Let $(\sigma_n, n\in\mathbb{N})$ be a sequence of positive integers satisfying $\sigma_n \to \infty$ as $n\to\infty$. Let b_n be uniformly distributed in $\mathfrak{B}_{\sigma_n}$, and let b_∞ be a uniform infinite bridge as specified in Section 4. Then, if k_n is a sequence of positive integers with $k_n \leqslant \sigma_n$ and $k_n = o(\sigma_n)$ as $n\to\infty$,*

$$\lim_{n\to\infty} \|\mathrm{Law}((\mathsf{b}_n(2\sigma_n-k_n),\dots,\mathsf{b}_n(2\sigma_n-1),\mathsf{b}_n(0),\mathsf{b}_n(1),\dots,\mathsf{b}_n(k_n))) \\ - \mathrm{Law}((\mathsf{b}_\infty(-k_n),\dots,\mathsf{b}_\infty(-1),\mathsf{b}_\infty(0),\mathsf{b}_\infty(1),\dots,\mathsf{b}_\infty(k_n)))\|_{\mathrm{TV}} = 0.$$

The proof follows from a small adaption of [8, Proof of Lemma 5.5] and is left to the reader. We stress, however, that in [8], b_n and b_∞ were defined in a slightly different manner, by grouping the +1-steps between two subsequent down-steps together to one "big" jump. Clearly, this does change the argument only in a minor way.

We are now in position to formulate an appropriate coupling of balls.

Proposition 4. *Fix $0<p<1/2$, and let $(\sigma_n, n\in\mathbb{N})$ be a sequence of positive integers satisfying $\sigma_n = \frac{1-2p}{p}n+o(n)$. Define γ_n in terms of σ_n as under (3), and put $\xi_n = \min\{n^{1/4}, \sqrt{n/\gamma_n}\}$. Then, given any $\varepsilon>0$, there exist $\delta>0$ and $n_0\in\mathbb{N}$ such that for every $n\geqslant n_0$, we can construct on the same probability space copies of $Q_n^{\sigma_n}$ and the UIHPQ_p such that with probability at least $1-\varepsilon$, the metric balls $B_{\delta\xi_n}(Q_n^{\sigma_n})$ and $B_{\delta\xi_n}(\mathsf{UIHPQ}_p)$ of radius $\delta\xi_n$ around the roots in the corresponding spaces are isometric.*

The local convergence of $Q_n^{\sigma_n}$ towards $\mathsf{UIHPQ}^{(p)}$ is a weaker statement, hence Theorem 1 in the case $0<p<1/2$ will follow from the proposition.

Proof. The proof is in spirit of [8, Proof of Proposition 3.11], requires, however, some modifications. We will indicate at which place we may simply adopt the reasoning. We consider a random uniform element $((\mathsf{f}_n,\mathsf{l}_n),\mathsf{b}_n)\in\mathfrak{F}^n_{\sigma_n}$, and a triplet $((\mathsf{f}_\infty^{(p)},\mathsf{l}_\infty^{(p)}),\mathsf{b}_\infty)$ consisting of a uniformly labeled infinite p-forest together with an (independent) uniform infinite bridge b_∞. We let $(Q_n^{\sigma_n},v^\bullet) = \Phi_n((\mathsf{f}_n,\mathsf{l}_n),\mathsf{b}_n)$ and $Q_\infty^\infty(p) = \Phi((\mathsf{f}_\infty^{(p)},\mathsf{l}_\infty^{(p)}),\mathsf{b}_\infty)$ be the quadrangulations obtained from applying the Bouttier-Di Francesco-Guitter mapping to $((\mathsf{f}_n,\mathsf{l}_n),\mathsf{b}_n)$ and $((\mathsf{f}_\infty^{(p)},\mathsf{l}_\infty^{(p)}),\mathsf{b}_\infty)$, respectively. Recall that $\mathsf{f}_n = (\mathsf{t}_0,\dots,\mathsf{t}_{\sigma_n-1})$ consists of σ_n trees. For $0\leqslant k\leqslant\sigma_n-1$, we let $\mathsf{t}(\mathsf{f}_n,k)=\mathsf{t}_k$, i.e., $\mathsf{t}(\mathsf{f}_n,k)$ is the tree of f_n with index k, and we put $\mathsf{t}(\mathsf{f}_n,\sigma_n)=\mathsf{t}(\mathsf{f}_n,0)$. In a similar manner, $\mathsf{t}(\mathsf{f}_\infty^{(p)},k)$ denotes the tree of $\mathsf{f}_\infty^{(p)}$ indexed by $k\in\mathbb{Z}$.

By Lemma 1, we find $\delta'>0$ and $n_0'\in\mathbb{N}$ such that for $n\geqslant n_0'$, we can construct $((\mathsf{f}_n,\mathsf{l}_n),\mathsf{b}_n)$ and $((\mathsf{f}_\infty^{(p)},\mathsf{l}_\infty^{(p)}),\mathsf{b}_\infty)$ on the same probability space such that with $A_n=\lfloor\delta'\xi_n^2\rfloor$, the event

$$\mathcal{E}^1(n,\delta') = \Big\{\mathsf{t}(\mathsf{f}_n,i)=\mathsf{t}(\mathsf{f}_\infty^{(p)},i),\ \mathsf{t}(\mathsf{f}_n,\sigma_n-i)=\mathsf{t}(\mathsf{f}_\infty^{(p)},-i) \text{ for all } 0\leqslant i\leqslant A_n\Big\} \\ \cap\Big\{\mathsf{l}_n|_{\mathsf{t}(\mathsf{f}_n,i)} = \mathsf{l}_\infty^{(p)}|_{\mathsf{t}(\mathsf{f}_\infty^{(p)},i)},\ \mathsf{l}_n|_{\mathsf{t}(\mathsf{f}_n,\sigma_n-i)} = \mathsf{l}_\infty^{(p)}|_{\mathsf{t}(\mathsf{f}_\infty^{(p)},-i)} \text{ for all } 0\leqslant i\leqslant A_n\Big\}$$

has probability at least $1-\varepsilon/8$. We now fix such a δ' for the rest of the proof. Recall that by our construction of the Bouttier-Di Francesco-Guitter bijection, the trees of $\mathfrak{f}_n$ are attached to the down-steps $\mathrm{d}_n^{\downarrow}(i) = \mathrm{d}_{\mathsf{b}_n}^{\downarrow}(i)$ of b_n, $1 \leqslant i \leqslant \sigma_n$, and similarly, the trees of $\mathfrak{f}_\infty^{(p)}$ are attached to the down-steps $\mathrm{d}_\infty^{\downarrow}(i) = \mathrm{d}_{\mathsf{b}_\infty}^{\downarrow}(i)$ of b_∞, where now $i \in \mathbb{Z}$. In view of the above event, this incites us to consider additionally the event

$$\begin{aligned}\mathcal{E}^2(n,\delta') = \quad & \left\{\mathsf{b}_n(i) = \mathsf{b}_\infty(i) \text{ for all } 1 \leqslant i \leqslant \mathrm{d}_\infty^{\downarrow}(A_n+1)\right\} \\ & \cap \left\{\mathsf{b}_n(2\sigma_n - i) = \mathsf{b}_\infty(i) \text{ for all } \mathrm{d}_\infty^{\downarrow}(-A_n) \leqslant i \leqslant -1\right\}.\end{aligned}$$

Note that on $\mathcal{E}^2(n,\delta')$, we automatically have $\mathrm{d}_n^{\downarrow}(i) = \mathrm{d}_\infty^{\downarrow}(i)$ for $1 \leqslant i \leqslant A_n + 1$, and $\mathrm{d}_n^{\downarrow}(\sigma_n - i + 1) = \mathrm{d}_\infty^{\downarrow}(-i)$ for $1 \leqslant i \leqslant A_n$. Trivially, we have that $\mathrm{d}_\infty^{\downarrow}(A_n+1) \geqslant A_n + 1$ and $\mathrm{d}_\infty^{\downarrow}(-A_n) \leqslant -A_n$, but also, with probability tending to 1, $\mathrm{d}_\infty^{\downarrow}(A_n+1) \leqslant 3A_n$ and $\mathrm{d}_\infty^{\downarrow}(-A_n) \geqslant -3A_n$. Since, in any case, $A_n = o(\sigma_n)$, we can ensure by Lemma 2 that the event $\mathcal{E}^2(n,\delta')$ has probability at least $1-\varepsilon/8$ for large n.

Now for $\delta > 0$, $n \in \mathbb{N}$, define the events

$$\mathcal{E}^3(n,\delta) = \left\{ \min_{[0,\, \mathrm{d}_\infty^{\downarrow}(A_n+1)]} \mathsf{b}_\infty < -5\delta\xi_n, \min_{[\mathrm{d}_\infty^{\downarrow}(-A_n),-1]} \mathsf{b}_\infty < -5\delta\xi_n \right\},$$

$$\mathcal{E}^4(n,\delta) = \left\{ \min_{[\mathrm{d}_\infty^{\downarrow}(A_n+1)+1,\, \mathrm{d}_\infty^{\downarrow}(-A_n)-1]} \mathsf{b}_n < -5\delta\xi_n \right\}.$$

By invoking Donsker's invariance principle together with Lemma 2 for the event $\mathcal{E}^3$ (and again the fact that $A_n + 1 \leqslant \mathrm{d}_\infty^{\downarrow}(A_n+1) \leqslant 3A_n$ and $-3A_n \leqslant \mathrm{d}_\infty^{\downarrow}(-A_n) \leqslant -A_n$ with high probability), we deduce that for small $\delta > 0$, provided n is large enough,

$$\mathbb{P}\left(\mathcal{E}^3(n,\delta)\right) \geqslant 1-\varepsilon/8, \quad \text{and} \quad \mathbb{P}\left(\mathcal{E}^4(n,\delta)\right) \geqslant 1-\varepsilon/8.$$

We will now assume that $n_0 \geqslant n_0'$ and $\delta > 0$ are such that for all $n \geqslant n_0$, the above bounds hold true, and work on the event $\mathcal{E}^1(n,\delta') \cap \mathcal{E}^2(n,\delta') \cap \mathcal{E}^3(n,\delta) \cap \mathcal{E}^4(n,\delta)$ of probability at least $1-\varepsilon/2$. We consider the forest obtained from restricting $\mathfrak{f}_n$ to the first A_n+1 and the last A_n trees,

$$\mathfrak{f}_n' = (\mathsf{t}(\mathfrak{f}_n, 0), \ldots, \mathsf{t}(\mathfrak{f}_n, A_n), \mathsf{t}(\mathfrak{f}_n, \sigma_n - A_n), \ldots, \mathsf{t}(\mathfrak{f}_n, \sigma_n - 1)).$$

Similarly, we define $\mathfrak{f}_\infty^{\prime(p)}$. We recall the cactus bounds in the version stated in [8, (4.4) of Section 4.5]. Applied to $Q_n^{\sigma_n}$, it shows that for vertices $v \in V(\mathfrak{f}_n)\backslash V(\mathfrak{f}_n')$, with d_n denoting the graph distance,

$$d_n(\mathfrak{f}_n(0), v) \geqslant -\max\left\{ \min_{[0,\, \mathrm{d}_\infty^{\downarrow}(A_n+1)]} \mathsf{b}_n, \min_{[\mathrm{d}_\infty^{\downarrow}(-A_n),\, 2\sigma_n]} \mathsf{b}_n \right\} \geqslant 5\delta\xi_n.$$

Applying now the analogous cactus bound [8, (4.6) of Section 4.5] to the infinite quadrangulation $Q_\infty^\infty(p)$, we obtain the same lower bound for vertices $v \in V(\mathfrak{f}_\infty^{(p)})\backslash V(\mathfrak{f}_\infty^{\prime(p)})$, with d_n replaced by the graph distance $d_\infty^{(p)}$ in $Q_\infty^\infty(p)$, and $\mathfrak{f}_n(0)$ replaced by the vertex $\mathfrak{f}_\infty^{(p)}(0)$ of $Q_\infty^\infty(p)$. We recall the definition of the metric balls $B_r^{(0)}(Q_n^{\sigma_n})$ and $B_r^{(0)}(Q_\infty^\infty(p))$; see Remark 9.

With the same arguments as in [8, Proof of Proposition 3.11], we then deduce that vertices at a distance at most $5\delta\xi_n - 1$ from $\mathfrak{f}_n(0)$ in $Q_n^{\sigma_n}$ agree with those at a distance at most $5\delta\xi_n - 1$ from $\mathfrak{f}_\infty^{(p)}(0)$ in $Q_\infty^\infty(p)$. Moreover,

$$d_n(u,v) = d_\infty^{(p)}(u,v) \quad \text{whenever } u, v \in B_{2\delta\xi_n}^{(0)}(Q_n^{\sigma_n}).$$

This proves that the balls $B_{2\delta\xi_n}^{(0)}(Q_n^{\sigma_n})$ and $B_{2\delta\xi_n}^{(0)}(Q_\infty^\infty(p))$ are isometric on an event of probability at least $1-\varepsilon/2$. In order to conclude, it suffices to observe that the distances from $\mathfrak{f}_n(0)$ resp. $\mathfrak{f}_\infty^{(p)}(0)$ to the root vertex in $Q_n^{\sigma_n}$ resp. $Q_\infty^\infty(p)$ are stochastically bounded; see again Remark 9. Clearly, this implies that with probability tending to 1 as n increases, we have the inclusions $B_{\delta\xi_n}(Q_n^{\sigma_n}) \subset B_{2\delta\xi_n}^{(0)}(Q_n^{\sigma_n})$ and $B_{\delta\xi_n}(Q_\infty^\infty(p)) \subset B_{2\delta\xi_n}^{(0)}(Q_\infty^\infty(p))$. □

As mentioned at the beginning, the case $p = 1/2$ has already been treated in [8, Proof of Proposition 3.11]: It is proved there that for δ small, balls of radius $\delta \min\{\sqrt{\sigma_n}, \sqrt{n/\sigma_n}\}$ in $Q_n^{\sigma_n}$ and in the standard $\mathsf{UIHPQ} = \mathsf{UIHPQ}_{1/2}$ can be coupled with high probability, implying of course again local convergence of $Q_n^{\sigma_n}$ towards the UIHPQ.

Finally, it remains to consider the case $p = 0$ corresponding to $\sigma_n \gg n$. This case is easy. We have the following coupling lemma.

Lemma 3. *Let $(\sigma_n, n \in \mathbb{N})$ be a sequence of positive integers satisfying $\sigma_n \gg n$. Put $\xi_n = \sigma_n/n$. Then, given any $\varepsilon > 0$, there exist $\delta > 0$ and $n_0 \in \mathbb{N}$ such that for every $n \geqslant n_0$, we can construct on the same probability space copies of $Q_n^{\sigma_n}$ and the UIHPQ_0 such that with probability at least $1-\varepsilon$, the metric balls $B_{\delta\xi_n}(Q_n^{\sigma_n})$ and $B_{\delta\xi_n}(\mathsf{UIHPQ}_0)$ of radius $\delta\xi_n$ around the roots in the corresponding spaces are isometric.*

Proof. Let $((\mathfrak{f}_n, \mathfrak{l}_n), \mathsf{b}_n) \in \mathfrak{F}_{\sigma_n}^n \times \mathfrak{B}_{\sigma_n}$ be uniformly distributed. By exchangeability of the trees, it follows that if $k_n = o(\sigma_n/n)$, then the first and last k_n trees of $\mathfrak{f}_n$ are all singletons with a probability tending to one. Applying Lemma 2, we can ensure that the event

$$\{\mathsf{b}_n(i) = \mathsf{b}_\infty(i),\ \mathsf{b}_n(2\sigma_n - i) = \mathsf{b}_\infty(-i),\ 1 \leqslant i \leqslant k_n\}$$

has a probability as large as we wish, provided n is large enough. Given $\varepsilon > 0$, the same arguments as in the proof of Proposition 4 yield an equality of balls $B_{\delta\xi_n}(Q_n^{\sigma_n})$ and $B_{\delta\xi_n}(\mathsf{UIHPQ}_0)$ for δ small and n large enough, on an event of probability at least $1-\varepsilon$. □

Let us now show that the space UIHPQ_0 defined in terms of the Bouttier-Di Francesco-Guitter mapping in Section 4.3 is nothing else than Kesten's tree associated to the critical geometric offspring law $\mu_{1/2}$.

Proof of Proposition 1. Let $\mathsf{b}_\infty = (\mathsf{b}_\infty(i), i \in \mathbb{Z})$ be a uniform infinite bridge, and let $(\mathfrak{f}_\infty^{(0)}, \mathfrak{l}_\infty^{(0)})$ be the infinite forest where all trees are just singletons (with label 0); see Section 4.3. The UIHPQ_0 is distributed as the infinite map $Q_\infty^\infty(0) = \Phi((\mathfrak{f}_\infty^{(0)}, \mathfrak{l}_\infty^{(0)}), \mathsf{b}_\infty)$. Since every vertex in $\mathfrak{f}_\infty^{(0)}$ defines a single corner, properties of the Bouttier-Di Francesco-Guitter mapping (Section 4.2) imply that $Q_\infty^\infty(0)$ is a tree almost surely. Moreover, the set of vertices of $Q_\infty^\infty(0)$ is identified with the set of down-steps $\mathsf{DS}(\mathsf{b}_\infty)$ of the bridge. Following [9, Section 2.2.3], conditionally on b_∞, we introduce a function $\varphi : \mathbb{Z} \to \mathsf{DS}(\mathsf{b}_\infty)$ that associates to $i \in \mathbb{Z}$ the next down-step $\geqslant i$ with label $\mathsf{b}_\infty(i)$ (and i is mapped to itself if $i \in \mathsf{DS}(\mathsf{b}_\infty)$). According to our rooting

convention, the root edge of $Q_\infty^\infty(0)$ connects $\varphi(0)$ to $\varphi(1)$. Note that φ is not injective almost surely.

We recall that Kesten's tree can be represented by a half-line of vertices $s_0, s_1, \dots$, together with a collection of independent Galton-Watson trees with offspring law $\mu_{1/2}$ grafted to the left and right side of each vertex s_i, $i \in \mathbb{N}_0$. We will now argue that the UIHPQ_0 $Q_\infty^\infty(0)$ has the same structure. In this regard, let us introduce the stopping times

$$S_i = \inf\{k \in \mathbb{N}_0 : \mathsf{b}_\infty(k) = -i\}, \quad i \in \mathbb{N}_0,$$

and denote by s_i the vertex of $Q_\infty^\infty(0)$ given by $\varphi(S_i)$. Together with their connecting edges, the collection $(s_i, i \in \mathbb{N}_0)$ forms a spine (i.e., an infinite self-avoiding path) in $Q_\infty^\infty(0)$.

The subtree rooted at s_i on the left side of the spine is encoded by the excursion $\{\mathsf{b}_\infty(k) : S_i \leqslant k \leqslant S_{i+1}\}$, in a way we describe next; see Figure 9 for an illustration. First note that by the Markov property, these subtrees for $i \in \mathbb{Z}$ are i.i.d.. In order to determine their law, let us consider the subtree encoded by the excursion $\{\mathsf{b}_\infty(k) : 0 \leqslant k \leqslant S_1\}$ of b_∞. This subtree is rooted at $s_0 = \varnothing$, and the number of offspring of s_0 is the number of down-steps with label 1 between 0 and S_1. Otherwise said, this is the number $\#\{0 < k < S_1 : \mathsf{b}_\infty(k) = 0\}$ of excursions of b_∞ above 0 between 0 and S_1. By the Markov property, this quantity follows the geometric distribution $\mu_{1/2}$ of parameter $1/2$. One can now repeat the argument for each child of s_0, by considering the corresponding excursion above 0 encoding its progeny tree, inside the mother excursion. We obtain that the subtree stemming from s_0 on the left of the spine has indeed the law of a Galton-Watson tree with offspring distribution $\mu_{1/2}$.

The subtrees attached to the vertices s_i, $i \in \mathbb{N}_0$, on the right of the spine can be treated by a symmetry argument. Namely, letting

$$S'_i = \inf\{k \in \mathbb{N}_0 : \mathsf{b}_\infty(-k) = -i\}, \quad i \in \mathbb{N}_0,$$

we observe that the subtree rooted at s_i to the right of the spine is coded by the (reversed) excursion $\{\mathsf{b}_\infty(k) : -S'_{i+1} \leqslant k \leqslant -S'_i\}$. With the same argument as above, we see that it has the law of an (independent) $\mu_{1/2}$-Galton Watson tree. This concludes the proof. □



Figure 9: The construction of the UIHPQ_0 from a uniform infinite bridge b_∞. The spine is shown in bold red arcs. The trees on the left of the spine are drawn in blue and enclosed by dotted blue half-circles, which indicate the corresponding excursions of b_∞ encoding these trees. The trees on the right of the spine are drawn in red, as the spine itself.

5.2 The UIHPQ_p as a local limit of Boltzmann quadrangulations

This section is devoted to the proof of Proposition 2. It is convenient to first prove the analogous result for pointed maps. For that purpose, we first extend the definitions of Boltzmann measures from Section 1.2.5 to pointed maps and then use a "de-pointing" argument. We use the notation $\mathcal{Q}_f^\bullet$ for the set of finite rooted pointed quadrangulations, and we write $\mathcal{Q}_f^{\bullet,\sigma}$ for the set of finite pointed rooted quadrangulations with 2σ boundary edges. The corresponding partition functions read

$$F^\bullet(g,z) = \sum_{\mathsf{q}\in\mathcal{Q}_f^\bullet} g^{\#\mathrm{F}(\mathsf{q})} z^{\#\partial\mathsf{q}/2}, \quad F_\sigma^\bullet(g) = \sum_{\mathsf{q}\in\mathcal{Q}_f^{\bullet,\sigma}} g^{\#\mathrm{F}(\mathsf{q})},$$

and the associated pointed Boltzmann distributions are defined by

$$\mathbb{P}_{g,z}^\bullet(\mathsf{q}) = \frac{g^{\#\mathrm{F}(\mathsf{q})} z^{\#\partial\mathsf{q}/2}}{F^\bullet(g,z)}, \quad \mathsf{q}\in\mathcal{Q}_f^\bullet, \quad \mathbb{P}_g^{\bullet,\sigma}(\mathsf{q}) = \frac{g^{\#\mathrm{F}(\mathsf{q})}}{F_\sigma^\bullet(g)}, \quad \mathsf{q}\in\mathcal{Q}_f^{\bullet,\sigma}.$$

We will need the following enumeration result for pointed rooted maps. From [16, (23)] and [15, Section 3.3], we have for every $0\leqslant p\leqslant 1/2$

$$F_\sigma^\bullet(g_p) = \binom{2\sigma}{\sigma}\left(\frac{1}{1-p}\right)^\sigma, \quad \sigma\in\mathbb{N}_0. \tag{7}$$

Note that the result (3.29) in [15] cannot be used directly, due to a difference in the rooting convention (there, the root vertex has to be chosen among the vertices of the boundary that are closest to the marked point).

Recall that $g_p = p(1-p)/3$ for $0\leqslant p\leqslant 1/2$. The first step towards the proof of Proposition 2 is the following convergence result for pointed Boltzmann quadrangulations.

Proposition 5. *Let $0\leqslant p\leqslant 1/2$. For every $\sigma\in\mathbb{N}_0$, let $Q_\sigma^\bullet(p)$ be a random rooted pointed quadrangulation distributed according to $\mathbb{P}_{g_p}^{\bullet,\sigma}$. Then, we have the local convergence for the metric d_{map} as $\sigma\to\infty$*

$$Q_\sigma^\bullet(p) \xrightarrow{(d)} \mathsf{UIHPQ}_p,$$

Proof. Let $\mathsf{q}\in\mathcal{Q}_f^\sigma$, and $((\mathfrak{f},\mathfrak{l}),\mathsf{b})\in\cup_{n\geqslant 0}\mathfrak{F}_\sigma^n\times\mathfrak{B}_\sigma$ such that $\mathsf{q}=\Phi((\mathfrak{f},\mathfrak{l}),\mathsf{b})$. Moreover, let $(\mathfrak{f}_\sigma^{(p)},\mathfrak{l}_\sigma^{(p)})$ be a uniformly labeled p-forest with σ trees, i.e., a collection of σ independent uniformly labeled p-Galton-Watson trees, and let b_σ be uniformly distributed in $\mathfrak{B}_\sigma$ and independent of $(\mathfrak{f}_\sigma^{(p)},\mathfrak{l}_\sigma^{(p)})$. We have

$$\begin{aligned}\mathbb{P}\left(\Phi\big((\mathfrak{f}_\sigma^{(p)},\mathfrak{l}_\sigma^{(p)}),\mathsf{b}_\sigma\big)=\mathsf{q}\right) &= \mathbb{P}\left(\big((\mathfrak{f}_\sigma^{(p)},\mathfrak{l}_\sigma^{(p)}),\mathsf{b}_\sigma\big)=((\mathfrak{f},\mathfrak{l}),\mathsf{b})\right)\\ &= \left(\frac{p(1-p)}{3}\right)^{|\mathfrak{f}|}\frac{(1-p)^\sigma}{\binom{2\sigma}{\sigma}} = \frac{g_p^{\#\mathrm{F}(\mathsf{q})}}{F_\sigma^\bullet(g_p)},\end{aligned}$$

Here, for the first equality in the second line, we have used (2), the fact that the label differences are i.i.d. uniform in $\{-1,0,1\}$, and $|\mathfrak{B}_\sigma|=\binom{2\sigma}{\sigma}$. The last equality follows from the enumeration result (7) and the fact that the number of edges of $\mathfrak{f}$ equals the number of faces of q. Thus, $Q_\sigma^\bullet(p)$ is distributed as $\Phi((\mathfrak{f}_\sigma^{(p)},\mathfrak{l}_\sigma^{(p)}),\mathsf{b}_\sigma)$.

Now observe that $\mathfrak{f}_\sigma^{(p)}$ is already a collection of σ independent p-Galton-Watson trees, and Lemma 2 allows us to couple the first and last $o(\sigma)$ steps of b_σ with those of a uniform infinite bridge b_∞. With exactly the same reasoning as in Proposition 4, we therefore obtain with high probability an isometry of balls $B_{\delta\sqrt{\sigma}}(Q_\sigma^\bullet(p))$ and $B_{\delta\sqrt{\sigma}}(\mathsf{UIHPQ}_p)$ for all σ sufficiently large, provided δ is small enough. The stated local convergence follows. □

Proposition 2 is a consequence of the foregoing result and the following de-pointing argument inspired by [1, Proposition 14]. According to Remark 2, it suffices to consider the case $p \in [0, 1/2)$.

In the following, by a small abuse of notation, we interpret $\mathbb{P}_{g_p}^{\bullet,\sigma}$ as a probability measure on $\mathcal{Q}_f$ by simply forgetting the marked point.

Lemma 4. *Let $0 \leqslant p < 1/2$. Then,*

$$\lim_{\sigma\to\infty} \left\| \mathbb{P}_{g_p}^{\sigma} - \mathbb{P}_{g_p}^{\bullet,\sigma} \right\|_{\mathrm{TV}} = 0.$$

Proof. Let $\#V$ be the mapping $\mathsf{q} \mapsto \#V(\mathsf{q})$, which assigns to a finite quadrangulation q its number of vertices. We have the absolute continuity relation [12, (5)]

$$\mathrm{d}\mathbb{P}_{g_p}^{\sigma}(\mathsf{q}) = \frac{K_\sigma}{\#V(\mathsf{q})} \mathrm{d}\mathbb{P}_{g_p}^{\bullet,\sigma}(\mathsf{q}),$$

where $K_\sigma = (\mathbb{E}_{g_p}^{\bullet,\sigma}[1/\#V)])^{-1}$. Then,

$$\left\| \mathbb{P}_{g_p}^{\sigma} - \mathbb{P}_{g_p}^{\bullet,\sigma} \right\|_{\mathrm{TV}} = \frac{1}{2} \sup_{F:\mathcal{Q}_f^\sigma \to [-1,1]} \left| \mathbb{E}_{g_p}^{\sigma}[F] - \mathbb{E}_{g_p}^{\bullet,\sigma}[F] \right| \leqslant \mathbb{E}_{g_p}^{\bullet,\sigma}\left[\left| 1 - \frac{K_\sigma}{\#V} \right| \right]. \tag{8}$$

Let $(\mathsf{t}_0^{(p)}, \dots, \mathsf{t}_{\sigma-1}^{(p)})$ be a collection of independent p-Galton-Watson trees. The proof of Proposition 5 shows that under $\mathbb{P}_{g_p}^{\bullet,\sigma}$, $\#V$ has the same law as

$$1 + \sum_{i=0}^{\sigma-1} \#V(\mathsf{t}_i^{(p)}).$$

Note that the summand $+1$ accounts for the pointed vertex, which is added to the tree vertices in the Bouttier-Di Francesco-Guitter mapping. Using the fact that $\#V(\mathsf{t}_0^{(p)})$ has the same law as $(T_{-1}^{(p)} + 1)/2$, where $T_{-1}^{(p)}$ is the first hitting time of -1 of a random walk with step distribution $p\delta_1 + (1-p)\delta_{-1}$, an application of the optional stopping theorem gives

$$\mathbb{E}_{g_p}^{\bullet,\sigma}[\#V] = 1 + \sigma\mathbb{E}\left[\#V(\mathsf{t}_0^{(p)})\right] = 1 + \sigma\left(\frac{1-p}{1-2p}\right).$$

Moreover, using $p < 1/2$ and the description in terms of $T_{-1}^{(p)}$, it is readily checked that the random variable $\#V(\mathsf{t}_0^{(p)})$ has small exponential moments. Cramer's theorem thus ensures that for every $\delta > 0$, there exists a constant $C_\delta > 0$ such that

$$\mathbb{P}_{g_p}^{\bullet,\sigma}\left(\left| \#V - \mathbb{E}_{g_p}^{\bullet,\sigma}[\#V] \right| > \delta\sigma \right) \leqslant \exp(-C_\delta\sigma).$$

We now proceed similarly to [1, Lemma 16]. Let X_σ be distributed as $\#V/\mathbb{E}^{\bullet,\sigma}_{g_p}[\#V]$ under $\mathbb{P}^{\bullet,\sigma}_{g_p}$. Note that $X_\sigma^{-1} \leqslant \mathbb{E}^{\bullet,\sigma}_{g_p}[\#V]$ $\mathbb{P}^{\bullet,\sigma}_{g_p}$-a.s. since $\#V \geqslant 1$. Moreover, it is seen that $\{|X_\sigma^{-1} - 1| > \delta\} \subset \{|X_\sigma| < 1/2\} \cup \{|X_\sigma - 1| > \delta/2\}$. From these observations, we obtain

$$\begin{aligned}\mathbb{E}\left[\left|X_\sigma^{-1} - 1\right|\right] &\leqslant \delta + \mathbb{E}\left[\left|X_\sigma^{-1} - 1\right| \mathbf{1}_{\{|X_\sigma^{-1}-1|>\delta\}}\right] \\ &\leqslant \delta + \left(\mathbb{E}^{\bullet,\sigma}_{g_p}[\#V] + 1\right) \mathbb{P}\left(|X_\sigma - 1| > \frac{\delta}{2} \wedge \frac{1}{2}\right).\end{aligned}$$

The preceding two displays show that the expected number of vertices grows linearly in σ, and the probability on the right decays exponentially fast in σ. Since $\delta > 0$ was arbitrary, we deduce that $X_\sigma^{-1} \longrightarrow 1$ as $\sigma \to \infty$ in $\mathbb{L}^1$. Finally,

$$\mathbb{E}^{\bullet,\sigma}_{g_p}\left[\left|1 - \frac{K_\sigma}{\#V}\right|\right] = \mathbb{E}\left[\left|1 - \frac{X_\sigma^{-1}}{\mathbb{E}[X_\sigma^{-1}]}\right|\right] \leqslant \frac{1}{\mathbb{E}[X_\sigma^{-1}]}\left(\left|\mathbb{E}[X_\sigma^{-1}] - 1\right| + \mathbb{E}[|X_\sigma^{-1} - 1|]\right) \longrightarrow 0$$

as $\sigma \to \infty$, which concludes the proof by (8). □

5.3 The BHP_θ as a local scaling limit of the UIHPQ_p's

In this section, we prove Theorem 2. For the reminder, we fix a sequence $(a_n, n \in \mathbb{N})$ of positive reals tending to infinity and let $r > 0$ be given. Similarly to [8, Proof of Theorem 3.4], the main step is to establish an absolute continuity relation of balls around the roots of radius ra_n between the UIHPQ_p for $p \in (0, 1/2]$ and the $\mathsf{UIHPQ} = \mathsf{UIHPQ}_{1/2}$. To this aim, we compute the Radon-Nikodym derivative of the encoding contour function of the UIHPQ_p with respect to that of the UIHPQ on an interval of the form $[-sa_n^2, sa_n^2]$ for $s > 0$. From Theorem 3.8 of [8] we know that $a_n^{-1} \cdot \mathsf{UIHPQ} \to \mathsf{BHP}_0$ in distribution in the local Gromov-Hausdorff topology, jointly with a uniform convergence on compacts of (rescaled) contour and label functions. An application of Girsanov's theorem shows that the limiting Radon-Nikodym derivative turns the contour function of BHP_0 into the contour function of BHP_θ, which allows us to conclude.

In order to make these steps rigorous, we begin with some notation specific to this section. Let $f \in \mathcal{C}(\mathbb{R}, \mathbb{R})$ and $x \in \mathbb{R}$. We define the last (first) visit to x to the left (right) of 0,

$$U_x(f) = \inf\{t \leqslant 0 : f(t) = x\} \in [-\infty, 0], \quad T_x(f) = \inf\{t \geqslant 0 : f(t) = x\} \in [0, \infty].$$

We agree that $U_x(f) = -\infty$ if the set over which the infimum is taken is empty, and, similarly, $T_x(f) = \infty$ if the second set is empty. We will also apply U_x to functions in $\mathcal{C}((-\infty, 0], \mathbb{R})$, and T_x to functions in $\mathcal{C}([0, \infty), \mathbb{R})$.

If $f \in \mathcal{C}(\mathbb{R}, \mathbb{R})$ is the contour function of an infinite p-forest for some $p \in (0, 1/2]$ (or part of it defined on some interval), and if $x \in \mathbb{N}$, we use the notation

$$\mathbf{v}(f, x) = \frac{1}{2}\left(T_{-x}(f) - U_x(f) - 2x\right)$$

for the total number of edges of the $2x$ trees encoded by f along the interval $[U_x(f), T_{-x}(f)]$. We set $\mathbf{v}(f, x) = \infty$ if $U_x(f)$ or $T_{-x}(f)$ is unbounded.

Given $s > 0$, we put for $n \in \mathbb{N}$

$$s_n = \lfloor (3/2) s a_n^2 \rfloor.$$

Now let $p \in (0, 1/2]$. Throughout this section and as usual, we assume that $((\mathfrak{f}_\infty^{(p)}, \mathfrak{l}_\infty^{(p)}), \mathsf{b}_\infty)$ and $((\mathfrak{f}_\infty, \mathfrak{l}_\infty), \mathsf{b}_\infty)$ encode the UIHPQ_p $Q_\infty^\infty(p)$ and the standard UIHPQ Q_∞^∞, respectively (see Definition 5). We stress that since the skewness parameter p does not affect the law of the infinite bridge b_∞, we can and will use the same bridge in the construction of both $Q_\infty^\infty(p)$ and Q_∞^∞. We denote by $(C_\infty^{(p)}, \mathfrak{L}_\infty^{(p)})$ and $(C_\infty, \mathfrak{L}_\infty)$ the associated contour and label functions, viewed as elements in $\mathcal{C}(\mathbb{R}, \mathbb{R})$.

For understanding how the balls of radius ra_n for some $r > 0$ around the roots in $Q_\infty^\infty(p)$ and Q_∞^∞ are related to each other, we need to control the contour functions $C_\infty^{(p)}$ and C_∞ on $[U_{s_n}, T_{-s_n}]$ for a suitable choice of $s = s(r)$. In this regard, we first formulate an absolute continuity relation between the probability laws $\mathbb{P}_{s,n}^{(p)}$ and $\mathbb{P}_{s,n}$ on $\mathcal{C}(\mathbb{R}, \mathbb{R})$ defined as follows:

$$\begin{aligned}
\mathbb{P}_{n,s}^{(p)} &= \mathrm{Law}\left(\left(C_\infty^{(p)}(t \vee U_{s_n}(C_\infty^{(p)}) \wedge T_{-s_n}(C_\infty^{(p)})), t \in \mathbb{R}\right)\right), \\
\mathbb{P}_{n,s} &= \mathrm{Law}\left((C_\infty(t \vee U_{s_n}(C_\infty) \wedge T_{-s_n}(C_\infty)), t \in \mathbb{R})\right).
\end{aligned}$$

Lemma 5. *Let $p \in (0, 1)$ and $s > 0$. The laws $\mathbb{P}_{n,s}^{(p)}$ and $\mathbb{P}_{n,s}$ are absolutely continuous with respect to each other: For any $f \in \mathrm{supp}(\mathbb{P}_{n,s}^{(p)})(= \mathrm{supp}(\mathbb{P}_{n,s}))$, with s_n as above,*

$$\mathbb{P}_{n,s}^{(p)}(f) = (4p(1-p))^{\mathbf{v}(f,s_n)} (2(1-p))^{2s_n} \mathbb{P}_{n,s}(f).$$

Proof. By definition of $C_\infty^{(p)}$ and C_∞, each element $f \in \mathcal{C}(\mathbb{R}, \mathbb{R})$ in the support of $\mathbb{P}_{n,s}^{(p)}$ lies also in the support of $\mathbb{P}_{n,s}$ and vice versa (note that $p \notin \{0, 1\}$).

More specifically, for such an f supported by these laws, $\mathbb{P}_{n,s}^{(p)}(f)$ resp. $\mathbb{P}_{n,s}(f)$ is the probability of a particular realization of $2s_n$ independent p-Galton-Watson trees resp. $(1/2)$-Galton-Watson trees with $\mathbf{v}(f, s_n)$ tree edges in total. Therefore, by (2),

$$\mathbb{P}_{n,s}^{(p)}(f) = p^{\mathbf{v}(f,s_n)}(1-p)^{\mathbf{v}(f,s_n)}(1-p)^{2s_n}, \quad \text{and} \quad \mathbb{P}_{n,s}(f) = 2^{-2(\mathbf{v}(f,s_n)+s_n)}.$$

This proves the lemma. □

We turn to the proof of Theorem 2. To that aim, we will work with rescaled and stopped versions of $(C_\infty^{(p)}, \mathfrak{L}_\infty^{(p)})$ and $(C_\infty, \mathfrak{L}_\infty)$, which encode the information of the first $s_n = \lfloor (3/2) s a_n^2 \rfloor$ trees to the right of zero, and of the first s_n trees to the left zero. Specifically, we let

$$\begin{aligned}
C_{n,s}^{\infty,p} &= \left(C_{n,s}^{\infty,p}(t), t \in \mathbb{R}\right) = \left(\frac{1}{(3/2)a_n^2} C_\infty^{(p)}\left((9/4)a_n^4 t \vee U_{s_n}(C_\infty^{(p)}) \wedge T_{-s_n}(C_\infty^{(p)})\right), t \in \mathbb{R}\right), \\
\mathfrak{L}_{n,s}^{\infty,p} &= \left(\mathfrak{L}_{n,s}^{\infty,p}(t), t \in \mathbb{R}\right) = \left(\frac{1}{a_n} \mathfrak{L}_\infty^{(p)}\left((9/4)a_n^4 t \vee U_{s_n}(C_\infty^{(p)}) \wedge T_{-s_n}(C_\infty^{(p)})\right), t \in \mathbb{R}\right), \\
C_{n,s}^{\infty} &= \left(C_{n,s}^{\infty}(t), t \in \mathbb{R}\right) = \left(\frac{1}{(3/2)a_n^2} C_\infty\left((9/4)a_n^4 t \vee U_{s_n}(C_\infty) \wedge T_{-s_n}(C_\infty)\right), t \in \mathbb{R}\right), \\
\mathfrak{L}_{n,s}^{\infty} &= \left(\mathfrak{L}_{n,s}^{\infty}(t), t \in \mathbb{R}\right) = \left(\frac{1}{a_n} \mathfrak{L}_\infty\left((9/4)a_n^4 t \vee U_{s_n}(C_\infty) \wedge T_{-s_n}(C_\infty)\right), t \in \mathbb{R}\right).
\end{aligned}$$

Following our notation from Section 3.1, we denote by $X^\theta = (X^\theta(t), t \in \mathbb{R})$ and $W^\theta = (W^\theta(t), t \in \mathbb{R})$ the contour and label functions of the limit space BHP_θ. We also put

$$X^{\theta,s} = \left(X^{\theta,s}(t), t \in \mathbb{R}\right) = \left(X^\theta\left(t \vee U_s(X^\theta) \wedge T_{-s}(X^\theta)\right), t \in \mathbb{R}\right),$$
$$W^{\theta,s} = \left(W^{\theta,s}(t), t \in \mathbb{R}\right) = \left(W^\theta\left(t \vee U_s(X^\theta) \wedge T_{-s}(X^\theta)\right), t \in \mathbb{R}\right).$$

Accordingly, we write X^0, W^0 and $X^{0,s}, W^{0,s}$ for the corresponding functions associated to BHP_0. We will make use of the following joint convergence.

Lemma 6. *Let $r, s > 0$. Then, in the notation from above, we have the joint convergence in law in $\mathcal{C}(\mathbb{R},\mathbb{R}) \times \mathcal{C}(\mathbb{R},\mathbb{R}) \times \mathbb{K}$,*

$$\left(C^\infty_{n,s}, \mathfrak{L}^\infty_{n,s}, B^{(0)}_r\left(a_n^{-1} \cdot Q^\infty_\infty\right)\right) \xrightarrow{(d)} \left(X^{0,s}, W^{0,s}, B_r(\mathsf{BHP}_0)\right).$$

Moreover, for $n \to \infty$

$$\frac{\mathbf{v}(C_\infty, s_n)}{(9/4)a_n^4} \xrightarrow{(d)} \frac{1}{2}\left(T_{-s} - U_s\right)(X^0).$$

Proof. Both statements are proved in [8]; to give a quick reminder, first note by standard random walk estimates that for each $\delta > 0$, there exists a constant $c_\delta > 0$ such that $\mathbb{P}(v(C_\infty, s_n) > c_\delta a_n^4) \leqslant \delta$; see [8, Proof of Lemma 6.18] for details. Together with the joint convergence in law in $\mathcal{C}(\mathbb{R},\mathbb{R})^2 \times \mathbb{K}$ obtained in [8, (6.30) of Remark 6.17], which reads

$$\left(\frac{C_\infty((9/4)a_n^4\cdot)}{(3/2)a_n^2}, \frac{\mathfrak{L}_\infty((9/4)a_n^4\cdot)}{a_n}, B^{(0)}_r\left(a_n^{-1} \cdot Q^\infty_\infty\right)\right) \xrightarrow{(d)} \left(X^0, W^0, B_r(\mathsf{BHP}_0)\right),$$

the first claim of the statement follows, and the second is then a consequence of this. □

We turn now to the Proof of Theorem 2.

Proof of Theorem 2. We fix a sequence $(p_n, n \in \mathbb{N}) \subset (0, 1/2]$ of the form

$$p_n = \frac{1}{2}\left(1 - \frac{2\theta}{3a_n^2}\right) + o\left(a_n^{-2}\right).$$

By Remark 9 and the observations in Section 1.2.7, the claim follows if we show that for all $r > 0$, as $n \to \infty$,

$$B^{(0)}_r\left(a_n^{-1} \cdot Q^\infty_\infty(p_n)\right) \xrightarrow{(d)} B_r(\mathsf{BHP}_\theta)$$

in distribution in $\mathbb{K}$. At this point, recall that $B^{(0)}_r(a_n^{-1} \cdot Q^\infty_\infty(p_n)) = a_n^{-1} \cdot B^{(0)}_{ra_n}(Q^\infty_\infty(p_n))$ is the (rescaled) ball of radius ra_n around the vertex $\mathfrak{f}^{(p_n)}_\infty(0)$ in $Q^\infty_\infty(p_n)$. We consider the event

$$\mathcal{E}^1(n,s) = \left\{\min_{[0,\,s_n]} \mathsf{b}_\infty < -3ra_n,\ \min_{[-s_n,\,0]} \mathsf{b}_\infty < -3ra_n\right\}.$$

We define a similar event in terms of the two-sided Brownian motion $\gamma = (\gamma(t), t \in \mathbb{R})$ scaled by the factor $\sqrt{3}$, which forms part of the construction of the space BHP_θ given in Section 3.1,

$$\mathcal{E}^2(s) = \left\{\min_{[0,s]} \gamma < -3r,\ \min_{[-s,0]} \gamma < -3r\right\}.$$

Using the cactus bound, it was argued in [8, Proof of Theorem 3.4] that on the event $\mathcal{E}^1(n,s)$, for any $p \in (0,1/2]$, the ball $B^{(0)}_{ra_n}(Q^\infty_\infty(p))$ viewed as a submap of $Q^\infty_\infty(p)$ is a measurable function of $(C^{\infty,p}_{n,s}, \mathfrak{L}^{\infty,p}_{n,s})$. (In [8], only the case $p = 1/2$ was considered, but the argument remains exactly the same for all p, since the encoding bridge b_∞ does not depend on the choice of p.) Similarly, on $\mathcal{E}^2(s)$, the ball $B_r(\mathsf{BHP}_\theta)$ for any $\theta \geqslant 0$ is a measurable function of $(X^{\theta,s}, W^{\theta,s})$.

Now let $\varepsilon > 0$ be given. By the (functional) central limit theorem, we find that for $s > 0$ and $n_0 \in \mathbb{N}$ sufficiently large, it holds that for all $n \geqslant n_0$, $\mathbb{P}(\mathcal{E}^1(n,s)) \geqslant 1-\varepsilon$. By choosing s possibly larger, we can moreover ensure that $\mathbb{P}(\mathcal{E}^2(s)) \geqslant 1-\varepsilon$. We fix such $s > 0$ and $n_0 \in \mathbb{N}$ such that for all $n \geqslant n_0$, both events $\mathcal{E}^1(n,s)$ and $\mathcal{E}^2(s)$ have probability at least $1-\varepsilon$.

Next, consider the laws $\mathbb{P}^{(p_n)}_{n,s}$ and $\mathbb{P}_{n,s}$ defined just above Lemma 5, and put for $f \in \mathcal{C}(\mathbb{R},\mathbb{R})$

$$\lambda_{n,s}(f) = (4p_n(1-p_n))^{\mathbf{v}(f,s_n)} \left(2(1-p_n)\right)^{2s_n}. \tag{9}$$

Then, with $F : \mathcal{C}(\mathbb{R},\mathbb{R})^2 \times \mathbb{K} \to \mathbb{R}$ measurable and bounded, Lemma 5 shows

$$\begin{aligned}\mathbb{E}\left[F\left(C^{\infty,p_n}_{n,s}, \mathfrak{L}^{\infty,p_n}_{n,s}, B^{(0)}_r\left(a_n^{-1}\cdot Q^\infty_\infty(p_n)\right)\right)\mathbf{1}_{\mathcal{E}^1(n,s)}\right] \\ = \mathbb{E}\left[\lambda_{n,s}(C_\infty) F\left(C^\infty_{n,s}, \mathfrak{L}^\infty_{n,s}, B^{(0)}_r\left(a_n^{-1}\cdot Q^\infty_\infty\right)\right)\mathbf{1}_{\mathcal{E}^1(n,s)}\right].\end{aligned} \tag{10}$$

Note that on the left side, we consider the closed ball of radius ra_n around the vertex $\mathfrak{f}_\infty(0)$ in the UIHPQ_{p_n} $Q^\infty_\infty(p_n)$, whereas on the right side, we look at the corresponding ball in the standard UIHPQ Q^∞_∞ with contour and label functions C_∞ and $\mathfrak{L}_\infty$. Plugging in the value of p_n in (9), we get

$$\lambda_{n,s}(f) = \left(1 + \frac{2\theta}{3a_n^2} + o(a_n^{-2})\right)^{2s_n} \left(1 - \frac{4\theta^2}{9a_n^4} + o(a_n^{-4})\right)^{\mathbf{v}(f,s_n)}. \tag{11}$$

Applying both statements of Lemma 6, and using (11), it follows that for large $n \geqslant n_1(\varepsilon)$

$$\begin{aligned}\Big|\mathbb{E}\left[\lambda_{n,s}(C_\infty) F\left(C^\infty_{n,s}, \mathfrak{L}^\infty_{n,s}, B^{(0)}_r\left(a_n^{-1}\cdot Q^\infty_\infty\right)\right)\right] - \\ \mathbb{E}\left[\exp\left(2s\theta - (T_{-s}-U_s)(X^0)\theta^2/2\right) F\left(X^{0,s}, W^{0,s}, B_r\left(\mathsf{BHP}_0\right)\right)\right]\Big| \leqslant \varepsilon.\end{aligned} \tag{12}$$

The rest of the proof is now similar to [8, Proof of Theorem 3.4]. Applying Pitman's transform and Girsanov's theorem, we have for continuous and bounded $G : \mathcal{C}(\mathbb{R},\mathbb{R})^2 \to \mathbb{R}$

$$\mathbb{E}\left[\exp\left(2s\theta - (T_{-s}-U_s)(X^0)\theta^2/2\right) G\left(X^{0,s}, W^{0,s}\right)\right] = \mathbb{E}\left[G\left(X^{\theta,s}, W^{\theta,s}\right)\right].$$

On $\mathcal{E}^2(s)$, $B_r(\mathsf{BHP}_0)$ is a measurable function of $(X^{0,s}, W^{0,s})$, and $B_r(\mathsf{BHP}_\theta)$ is given by the *same* measurable function of $(X^{\theta,s}, W^{\theta,s})$. Consequently,

$$\begin{aligned}\mathbb{E}\left[\exp\left(2s\theta - (T_{-s}-U_s)(X^0)\theta^2/2\right) F\left(X^{0,s}, W^{0,s}, B_r\left(\mathsf{BHP}_0\right)\right)\mathbf{1}_{\mathcal{E}^2(s)}\right] \\ = \mathbb{E}\left[F\left(X^{\theta,s}, W^{\theta,s}, B_r\left(\mathsf{BHP}_\theta\right)\right)\mathbf{1}_{\mathcal{E}^2(s)}\right].\end{aligned} \tag{13}$$

Recall that the events $\mathcal{E}^1(n,s)$ and $\mathcal{E}^2(s)$ have probability at least $1-\varepsilon$. Using this fact together with (10), (12), (13) and the triangle inequality, we find a constant $C = C(F,s,\theta)$ such that for sufficiently large n,

$$\left|\mathbb{E}\left[F\left(C^{\infty,p_n}_{n,s}, \mathfrak{L}^{\infty,p_n}_{n,s}, B^{(0)}_r\left(a_n^{-1}\cdot Q^\infty_\infty(p_n)\right)\right)\right] - \mathbb{E}\left[F\left(X^{\theta,s}, W^{\theta,s}, B_r\left(\mathsf{BHP}_\theta\right)\right)\right]\right| \leqslant C\varepsilon.$$

This implies the theorem. □

5.4 The ICRT as a local scaling limit of the UIHPQ$_p$'s

Theorem 3 states that the ICRT appears as the distributional limit of $a_n^{-1} \cdot \mathsf{UIHPQ}_{p_n}$ when $a_n \to \infty$ and $p_n \in [0, 1/2]$ satisfies $a_n^2 (1 - 2p_n) \to \infty$ as $n \to \infty$. In essence, the idea behind the proof is the following. Fix $r > 0$, and sequences $(a_n)_n$ and $(p_n)_n$ with the above properties. It turns out that in the UIHPQ_{p_n}, vertices at a distance less than ra_n from the root are to be found at a distance of order $o(a_n)$ from the boundary. Therefore, upon rescaling the graph distance by a factor a_n^{-1}, the scaling limit of the UIHPQ_{p_n} in the local Gromov-Hausdorff sense will agree with the scaling limit of its boundary. Upon a rescaling by a_n^2 in time and a_n^{-1} in space, the encoding bridge b_∞ converges to a two-sided Brownian motion, which in turn encodes the ICRT.

The above observations are most naturally turned into a proof using the description of the Gromov-Hausdorff metric in terms of correspondences between metric spaces; see [18, Theorem 7.3.25]. Lemma 10 below captures the kind of correspondence we need to construct. Our strategy of showing convergence of quadrangulations with a boundary towards a tree has already been successfully implemented before; see, for instance, [11, Proof of Theorem 5].

For the reminder of this section, we write $((\mathfrak{f}_\infty^{(n)}, \mathfrak{l}_\infty^{(n)}), \mathsf{b}_\infty)$ for a uniformly labeled infinite p_n-forest together with an (independent) uniform infinite bridge b_∞, and we assume that the UIHPQ_{p_n} is given in terms of $((\mathfrak{f}_\infty^{(n)}, \mathfrak{l}_\infty^{(n)}), \mathsf{b}_\infty)$, *via* the Bouttier-Di Francesco-Guitter mapping. We interpret the associated contour function $C_\infty^{(n)}$, the bridge b_∞ and the (unshifted) labels $\mathfrak{l}_\infty^{(n)}$ as elements in $\mathcal{C}(\mathbb{R}, \mathbb{R})$ (by linear interpolation); see Section 4.1.2.

The core of the argument lies in the following lemma, which gives the necessary control over distances to the boundary, *via* a control of the labels $\mathfrak{l}_\infty^{(n)}$. We will use it at the very end of the proof of Theorem 3, which follows afterwards.

Lemma 7. *Let $(a_n, n \in \mathbb{N})$ be a sequence of positive reals tending to infinity, and $(p_n, n \in \mathbb{N}) \subset [0, 1/2)$ be a sequence satisfying $a_n^2(1 - 2p_n) \to \infty$ as $n \to \infty$. Then, in the notation from above, we have the distributional convergence in $\mathcal{C}(\mathbb{R}, \mathbb{R}^2)$ as $n \to \infty$,*

$$\left(\left(\frac{1}{a_n^2} C_\infty^{(n)} \left(\frac{a_n^2}{1 - 2p_n} s \right), \frac{1}{a_n} \mathfrak{l}_\infty^{(n)} \left(\frac{a_n^2}{1 - 2p_n} s \right), s \in \mathbb{R} \right) \xrightarrow{(d)} ((-s, 0), s \in \mathbb{R}) \right).$$

Proof. We have to show joint convergence of $C_\infty^{(n)}$ and $\mathfrak{l}_\infty^{(n)}$ on any interval of the form $[-K, K]$, for $K > 0$. Due to an obvious symmetry in the definition of the contour function, we may restrict ourselves to intervals of the form $[0, K]$. Fix $K > 0$, and put $\theta_n = (1 - 2p_n)^{-1} a_n^2$. We first show that $a_n^{-2} C_\infty^{(n)}(\theta_n s)$, $s \in \mathbb{R}$, converges on $[0, K]$ to $g(s) = -s$ in probability. For that purpose, recall that $C_\infty^{(n)}$ on $[0, \infty)$ has the law of an linearly interpolated random walk started from 0 with step distribution $p_n \delta_1 + (1 - p_n)\delta_{-1}$. Set $K_n = \lceil K\theta_n \rceil$, and let $\delta > 0$. By using Doob's inequality in the second line,

$$\begin{aligned} \mathbb{P}\Big(\sup_{s \in [0,K]} \left| a_n^{-2} C_\infty^{(n)}(\theta_n s) + s \right| > \delta \Big) &\leqslant \mathbb{P}\Big(\sup_{0 \leqslant i \leqslant K_n} \left| C_\infty^{(n)}(i) + (1 - 2p_n) i \right| > \delta a_n^2 \Big) \\ &\leqslant \frac{1}{\delta^2 a_n^4} \mathbb{E}\left[\left| C_\infty^{(n)}(K_n) + (1 - 2p_n) K_n \right|^2 \right] \leqslant \frac{4K_n}{\delta^2 a_n^4} \leqslant \frac{4K}{\delta^2 a_n^2 (1 - 2p_n)}. \end{aligned} \tag{14}$$

Thanks to our assumption on p_n, the right hand side converges to zero, and the convergence of the contour function is established. Showing joint convergence together with the (rescaled) labels $\mathfrak{l}_\infty^{(n)}$ is now rather standard: First, we may assume by Skorokhod's theorem that $a_n^{-2}C_\infty^{(n)}(\theta_n s)$ converges on $[0,K]$ almost surely. Now fix $0 \leqslant s \leqslant K$. Conditionally given $C_\infty^{(n)}$ on $[0, K\theta_n]$, we have by construction, for $(\eta_i, i \in \mathbb{N})$ a sequence of i.i.d. uniform random variables on $\{-1,0,1\}$, and with $\underline{C}_\infty^{(n)}(\lfloor \theta_n s \rfloor) = \min_{[0,\lfloor \theta_n s \rfloor]} C_\infty^{(n)}$,

$$\mathfrak{l}_\infty^{(n)}(\lfloor \theta_n s \rfloor) =_d \sum_{i=1}^{C_\infty^{(n)}(\lfloor \theta_n s \rfloor) - \underline{C}_\infty^{(n)}(\lfloor \theta_n s \rfloor)} \eta_i. \tag{15}$$

Conditionally given $C_\infty^{(n)}$ on $[0, K\theta_n]$, for $\delta > 0$, Chebycheff's inequality gives

$$\mathbb{P}\left(\mathfrak{l}_\infty^{(n)}(\lfloor \theta_n s \rfloor) > \delta a_n \,|\, C_\infty^{(n)} \restriction [0, \theta_n s]\right) \leqslant \frac{1}{\delta^2 a_n^2}\left(C_\infty^{(n)}(\lfloor \theta_n s \rfloor) - \underline{C}_\infty^{(n)}(\lfloor \theta_n s \rfloor)\right).$$

By our assumption, $a_n^{-2}(C_\infty^{(n)}(\lfloor \theta_n s \rfloor) - \underline{C}_\infty^{(n)}(\lfloor \theta_n s \rfloor))$ converges to zero almost surely, and we conclude

$$\left(a_n^{-2}C_\infty^{(n)}(\lfloor \theta_n s \rfloor), a_n^{-1}\mathfrak{l}_\infty^{(n)}(\lfloor \theta_n s \rfloor)\right) \xrightarrow{(d)} (-s, 0) \quad \text{as } n \to \infty.$$

Since both $C_\infty^{(n)}$ and $\mathfrak{l}_\infty^{(n)}$ are Lipschitz almost surely, the claim follows with $\lfloor \theta_n s \rfloor$ replaced by $\theta_n s$. Joint finite-dimensional convergence can now be shown inductively: As for two-dimensional convergence on $[0,K]$, we simply note that when $0 \leqslant s_1 < s_2 \leqslant K$ are such that $C_\infty^{(n)}(\lfloor \theta_n s_1 \rfloor)$ and $C_\infty^{(n)}(\lfloor \theta_n s_2 \rfloor)$ encode vertices of different trees of $\mathfrak{f}_\infty^{(n)}$, then, conditionally on $C_\infty^{(n)} \restriction [0, \lfloor \theta_n s_2 \rfloor]$, $\mathfrak{l}_\infty^{(n)}(\lfloor \theta_n s_1 \rfloor)$ and $\mathfrak{l}_\infty^{(n)}(\lfloor \theta_n s_2 \rfloor)$ are independent sums of i.i.d. uniform variables on $\{-1,0,1\}$, and we have a representation similar to (15). If $C_\infty^{(n)}(\lfloor \theta_n s_1 \rfloor)$ and $C_\infty^{(n)}(\lfloor \theta_n s_2 \rfloor)$ encode vertices of one and the same tree of $\mathfrak{f}_\infty^{(n)}$, then, with the abbreviation

$$\check{C}_\infty^{(n)}(s_1, s_2) = \min_{[\lfloor \theta_n s_1 \rfloor, \lfloor \theta_n s_2 \rfloor]} C_\infty^{(n)} - \underline{C}_\infty^{(n)}(\lfloor \theta_n s_1 \rfloor),$$

it holds that

$$\begin{aligned}
\mathfrak{l}_\infty^{(n)}(\lfloor \theta_n s_1 \rfloor) &=_d \sum_{i=1}^{\check{C}_\infty^{(n)}(s_1,s_2)} \eta_i + \sum_{i=\check{C}_\infty^{(n)}(s_1,s_2)+1}^{C_\infty^{(n)}(\lfloor \theta_n s_1 \rfloor)} \eta_i', \\
\mathfrak{l}_\infty^{(n)}(\lfloor \theta_n s_2 \rfloor) &=_d \sum_{i=1}^{\check{C}_\infty^{(n)}(s_1,s_2)} \eta_i + \sum_{i=\check{C}_\infty^{(n)}(s_1,s_2)+1}^{C_\infty^{(n)}(\lfloor \theta_n s_2 \rfloor)} \eta_i',
\end{aligned}$$

where $(\eta_i', i \in \mathbb{N})$ is an i.i.d. copy of $(\eta_i, i \in \mathbb{N})$. Using almost sure convergence of $a_n^{-2}C_\infty^{(n)}(\theta_n s)$ on $[0,K]$ and an argument similar to that in the one-dimensional convergence considered above, we get two-dimensional convergence of $(a_n^{-2}C_\infty^{(n)}(\theta_n s), a_n^{-1}\mathfrak{l}_\infty^{(n)}(\theta_n s))$ on $[0,K]$, as wanted. Some more details can be found in [34, Proof of Theorem 4.3]. Higher-dimensional convergence is now shown inductively and is left to the reader. It remains to show tightness of the rescaled labels. We begin with the following lemma.

Lemma 8. *Let $K > 0$, $(a_n, n \in \mathbb{N})$ and $(p_n, n \in \mathbb{N})$ be as above. Then, for any $q \geqslant 2$, there exists a constant $C_q > 0$ such that for any $n \in \mathbb{N}$ and any $0 \leqslant s_1, s_2 \leqslant K$, we have (with $\theta_n = (1-2p_n)^{-1}a_n^2$, as before)*

$$a_n^{-2q}\,\mathbb{E}\left[\left|C_\infty^{(n)}(\theta_n s_1) - C_\infty^{(n)}(\theta_n s_2)\right|^q\right] \leqslant C_q|s_1 - s_2|^{q/2}.$$

Proof. If $|s_1 - s_2| \leqslant \theta_n^{-1}$, then, using linearity of $C_\infty^{(n)}$,

$$a_n^{-2q}\,\mathbb{E}\left[\left|C_\infty^{(n)}(\theta_n s_1) - C_\infty^{(n)}(\theta_n s_2)\right|^q\right] \leqslant a_n^{-2q}\theta_n^q|s_1 - s_2|^q \leqslant a_n^{-2q}\theta_n^{q/2}|s_1 - s_2|^{q/2}.$$

Since $a_n^{-2q}\theta_n^{q/2} \leqslant a_n^{-q}(1-2p_n)^{-q/2} \to 0$ by assumption on p_n, the claim of the lemma follows in this case. Now let $|s_1 - s_2| > \theta_n^{-1}$. We may assume $s_2 \geqslant s_1$. Using the triangle inequality and again the assumption on p_n, we see that it suffices to establish the claim in the case where $\theta_n s_1$ and $\theta_n s_2$ are integers. In this case, by definition of $C_\infty^{(n)}$,

$$C_\infty^{(n)}(\theta_n s_2) - C_\infty^{(n)}(\theta_n s_1) =_d \Big(\sum_{i=1}^{\theta_n(s_2-s_1)} \vartheta_i\Big) - \theta_n(s_2 - s_1)(1-2p_n),$$

where $(\vartheta_i, i \in \mathbb{N})$ are (centered) i.i.d. random variables with distribution $p_n\delta_{2(1-p_n)} + (1-p_n)\delta_{-2p_n}$. Using that $|a+b|^q \leqslant 2^{q-1}(|a|^q + |b|^q)$ for reals a, b, we get

$$\mathbb{E}\left[\left|C_\infty^{(n)}(\theta_n s_2) - C_\infty^{(n)}(\theta_n s_1)\right|^q\right] \leqslant 2^{q-1}\Big(\mathbb{E}\Big[\Big|\sum_{i=1}^{\theta_n(s_2-s_1)} \vartheta_i\Big|^q\Big] + \theta_n^q(1-2p_n)^q(s_2-s_1)^q\Big).$$

The second term within the parenthesis is equal to $a_n^{2q}|s_2 - s_1|^q \leqslant K^{q/2}a_n^{2q}|s_2 - s_1|^{q/2}$. As for the sum, we apply Rosenthal's inequality and obtain for some constant $C_q' > 0$,

$$\mathbb{E}\Big[\Big|\sum_{i=1}^{\theta_n(s_2-s_1)} \vartheta_i\Big|^q\Big] \leqslant C_q'\theta_n^{q/2}|s_2 - s_1|^{q/2}.$$

Using once more that $a_n^{-2q}\theta_n^{q/2} \to 0$ by assumption on p_n, the lemma is proved. □

Let $\kappa > 0$. By the theorem of Kolmogorov-Čentsov (see [29, Theorem 2.8]), it follows from the above lemma that there exists $M = M(\kappa) > 0$ such that for all $n \in \mathbb{N}$, the event

$$\mathcal{E}_n = \left\{\sup_{0\leqslant s<t\leqslant K} \frac{|C_\infty^{(n)}(\theta_n s) - C_\infty^{(n)}(\theta_n t)|}{a_n^2|s-t|^{2/5}} \leqslant M\right\}$$

has probability at least $1 - \kappa$. We will now work conditionally given $\mathcal{E}_n$.

Lemma 9. *In the setting from above, there exists a constant $C' > 0$ such that for all $n \in \mathbb{N}$ and all $0 \leqslant s_1, s_2 \leqslant K$,*

$$\mathbb{E}\left[a_n^{-6}|\mathfrak{l}_\infty^{(n)}(\theta_n s_1) - \mathfrak{l}_\infty^{(n)}(\theta_n s_2)|^6 \,\middle|\, \mathcal{E}_n\right] \leqslant C'|s_1 - s_2|^{6/5}.$$

Tightness of the conditional laws of $a_n^{-1}\mathfrak{l}_\infty^{(n)}(\theta_n s)$, $0 \leqslant s \leqslant K$, given $\mathcal{E}_n$ is a standard consequence of this lemma; see [29, Problem 4.11]). Since κ in the definition of $\mathcal{E}_n$ can be chosen arbitrarily small, tightness of the unconditioned laws of the rescaled labels follows, and so does Lemma 7. □

It therefore only remains to prove Lemma 9.

Proof of Lemma 9. With arguments similar to those in the proof of Lemma 8, we see that it suffices to prove the claim in the case where $\theta_n s_1$ and $\theta_n s_2$ are integers (and $s_1 \leqslant s_2$). Let

$$\Delta C_\infty^{(n)}(s_1, s_2) = C_\infty^{(n)}(\theta_n s_1) + C_\infty^{(n)}(\theta_n s_2) - 2 \min_{[\theta_n s_1, \theta_n s_2]} C_\infty^{(n)}.$$

By definition of $(C_\infty^{(n)}, \mathfrak{l}_\infty^{(n)})$, conditionally given $C_\infty^{(n)}$ on $[0, K]$, the difference $|\mathfrak{l}_\infty^{(n)}(\theta_n s_2) - \mathfrak{l}_\infty^{(n)}(\theta_n s_1)|$ is distributed as a sum of i.i.d. variables η_i with the uniform law on $\{-1, 0, 1\}$. By construction, the sum involves at most $\Delta C_\infty^{(n)}(s_1, s_2)$ summands: Indeed, it involves exactly $\Delta C_\infty^{(n)}(s_1, s_2)$ many summands if $C_\infty^{(n)}(\theta_n s_1)$ and $C_\infty^{(n)}(\theta_n s_2)$ encode vertices of the same tree, and less than $\Delta C_\infty^{(n)}(s_1, s_2)$ many summands if they encode vertices of different trees. Again with Rosenthal's inequality, we thus obtain for some $\tilde{C} > 0$,

$$\begin{aligned}\mathbb{E}\left[a_n^{-6}|\mathfrak{l}_\infty^{(n)}(\theta_n s_2) - \mathfrak{l}_\infty^{(n)}(\theta_n s_1)|^6 \,\middle|\, \mathcal{E}_n\right] &\leqslant a_n^{-6}\mathbb{E}\left[\left|\sum_{i=1}^{\Delta C_\infty^{(n)}(s_1,s_2)} \eta_i\right|^6 \,\middle|\, \mathcal{E}_n\right] \\ &\leqslant \tilde{C} a_n^{-6}\mathbb{E}\left[|\Delta C_\infty^n(s_1, s_2)|^3 \,\middle|\, \mathcal{E}_n\right].\end{aligned}$$

On $\mathcal{E}_n$, we have the bound

$$a_n^{-2}|\Delta C_\infty^n(s_1, s_2)| \leqslant 2 \sup_{0 \leqslant s < t \leqslant K} \frac{|C_\infty^n(\theta_n s) - C_\infty^n(\theta_n t)|}{a_n^2 |s - t|^{2/5}} |s_1 - s_2|^{2/5} \leqslant M|s_1 - s_2|^{2/5},$$

and the claim of the lemma follows. □

Finally, for proving Theorem 3, we will make use of the following lemma.

Lemma 10 (Lemma 5.7 of [8])**.** *Let $r \geqslant 0$. Let $\mathbf{E} = (E, d, \rho)$ and $\mathbf{E}' = (E', d', \rho')$ be two pointed complete and locally compact length spaces. Consider a subset $\mathcal{R} \subset E \times E'$ which has the following properties:*

- *$(\rho, \rho') \in \mathcal{R}$,*
- *for all $x \in B_r(\mathbf{E})$, there exists $x' \in E'$ such that $(x, x') \in \mathcal{R}$,*
- *for all $y' \in B_r(\mathbf{E}')$, there exists $y \in E$ such that $(y, y') \in \mathcal{R}$.*

Then, $d_{\mathrm{GH}}(B_r(\mathbf{E}), B_r(\mathbf{E}')) \leqslant (3/2) \sup\{|d(x, y) - d'(x', y')| : (x, x'), (y, y') \in \mathcal{R}\}$.

A proof is given in [8]. Although $\mathcal{R}$ is not necessarily a correspondence in the sense of [18], we might call the supremum on the right side of the inequality the *distortion* of $\mathcal{R}$.

Proof of Theorem 3. We let $(a_n, n \in \mathbb{N})$ and $(p_n, n \in \mathbb{N}) \subset [0, 1/2]$ be two sequences as in the statement, and, as mentioned at the beginning of this section, we assume that the UIHPQ_{p_n} $Q_\infty^\infty(p_n)$ with skewness parameter p_n is encoded in terms of $((\mathfrak{f}_\infty^{(n)}, \mathfrak{l}_\infty^{(n)}), \mathsf{b}_\infty)$. Local Gromov-Hausdorff convergence in law of $a_n^{-1} \cdot Q_\infty^\infty(p_n)$ towards the ICRT follows if we prove that for each $r \geqslant 0$,

$$B_r^{(0)}\left(a_n^{-1} \cdot Q_\infty^\infty(p_n)\right) \xrightarrow{(d)} B_r(\mathsf{ICRT}) \tag{16}$$

in distribution in $\mathbb{K}$, where we recall again that $B_r^{(0)}(a_n^{-1} \cdot Q_\infty^\infty(p_n))$ denotes the ball of radius r around the vertex $\mathfrak{f}_\infty^{(n)}(0)$ in the rescaled UIHPQ_{p_n}.

We will show the claim for $r = 1$. The proof follows essentially the line of argumentation in [8, Proof of Theorem 3.5]; since the argument is short, we repeat the main steps for completeness. We will apply Lemma 10 in the following way. The ICRT takes the role of the space $\mathbf{E}'$, with the equivalence class $[0]$ of zero being the distinguished point. Then, we consider for each $n \in \mathbb{N}$ the space $a_n^{-1} \cdot Q_\infty^\infty(p_n)$ pointed at $\mathfrak{f}_\infty^{(n)}(0)$, which takes the role of $\mathbf{E}$ in the lemma. We construct a subset $\mathcal{R}_n \subset E \times E'$ with the properties of Lemma 10, such that its distortion, that is, the quantity $\sup\{|d(x, y) - d'(x', y')| : (x, x'), (y, y') \in \mathcal{R}_n\}$, is of order $o(1)$ for n tending to infinity. By Lemma 10, this will prove (16). We remark that $Q_\infty^\infty(p_n)$ is not a length space, hence Lemma 10 seems not applicable at first sight. However, as explained in Section 1.2.7, by identifying each edge with a copy of $[0, 1]$ and upon extending the graph metric isometrically, we may identify $Q_\infty^\infty(p_n)$ with the (associated) length space, which we denote by $\mathbf{Q}_\infty^\infty(p_n) = (V(\mathbf{Q}_\infty^\infty(p_n)), d_{\mathrm{gr}}, \rho)$. Here and in what follows, $\boldsymbol{d}_{\mathrm{gr}}$ is the graph metric isometrically extended to $\mathbf{Q}_\infty^\infty(p_n)$. Note that the vertex set $V(\mathfrak{f}_\infty^{(n)})$ may be viewed as a subset of $\mathbf{Q}_\infty^\infty(p_n)$, and between points of $V(\mathfrak{f}_\infty^{(n)})$, the distances $\boldsymbol{d}_{\mathrm{gr}}$ and d_{gr} agree. Moreover, as a matter of fact, every point in $\mathbf{Q}_\infty^\infty(p_n)$ is at distance at most $1/2$ away from a vertex of $\mathfrak{f}_\infty^{(n)}$.

Recall that $(\mathsf{b}_\infty(t), t \in \mathbb{R})$ has the law of a (linearly interpolated) two-sided symmetric simple random walk with $\mathsf{b}_\infty(0) = 0$. Let $X = (X_t, t \in \mathbb{R})$ be a two-sided Brownian motion with $X_0 = 0$. By Donsker's invariance principle, we deduce that as n tends to infinity,

$$\left(a_n^{-1}\mathsf{b}_\infty(a_n^2 t), t \in \mathbb{R}\right) \xrightarrow{(d)} (X_t, t \in \mathbb{R})\,. \tag{17}$$

Using Skorokhod's representation theorem, we can assume that the above convergence holds almost surely on a common probability space, uniformly over compacts. Now let $\delta > 0$, and fix $\alpha > 0$ and $n_0 \in \mathbb{N}$ such that the event

$$\mathcal{E}(n, \alpha) = \left\{\max\left\{\min_{[0,\alpha]} X, \min_{[-\alpha,0]} X\right\} < -1\right\} \bigcap \left\{\max\left\{\min_{[0,\alpha a_n^2]} \mathsf{b}_\infty, \min_{[-\alpha a_n^2,0]} \mathsf{b}_\infty\right\} < -a_n\right\}$$

has probability at least $1 - \delta$ for $n \geqslant n_0$. From now on, we argue on the event $\mathcal{E}(n, \alpha)$. We moreover assume that the ICRT $(\mathcal{T}_X, d_X, [0])$ is defined in terms of X, and we write $p_X : \mathbb{R} \to \mathcal{T}_X$ for the canonical projection.

Recall that the vertices of $\mathfrak{f}_\infty^{(n)} = (\mathfrak{t}_i, i \in \mathbb{Z})$ are identified with the vertices of $Q_\infty^\infty(p_n)$. The mapping $\mathcal{I}(v) \in \mathbb{Z}$ gives back the index of the tree a vertex $v \in V(\mathfrak{f}_\infty^{(n)})$ belongs to. We extend $\mathcal{I}$ to the elements of the length space $\mathbf{Q}_\infty^\infty(p_n)$ as follows. By viewing $V(\mathfrak{f}_\infty^{(n)})$ as a subset of $\mathbf{Q}_\infty^\infty(p_n)$ as explained above, we associate to every point u of $\mathbf{Q}_\infty^\infty(p_n)$ its closest vertex $v \in V(\mathfrak{f}_\infty^{(n)})$ satisfying $\boldsymbol{d}_{\mathrm{gr}}(\mathfrak{f}_\infty^{(n)}(0), v) \geqslant d_{\mathrm{gr}}(\mathfrak{f}_\infty^{(n)}(0), u)$. Note again $\boldsymbol{d}_{\mathrm{gr}}(v, u) \leqslant 1/2$. Put

$$A_n = \{u \in \mathbf{Q}_\infty^\infty(p_n) : \mathcal{I}(u) \in [-\alpha a_n^2, \alpha a_n^2]\}.$$

A direct application of the cactus bound [8, (4.6) of Section 4.5] shows that on $\mathcal{E}(n,\alpha)$,

$$\boldsymbol{d}_{\mathrm{gr}}(\mathfrak{f}_\infty^{(n)}(0),u) > a_n \quad \text{whenever } \mathcal{I}(u) \notin A_n,$$

implying that the set A_n contains the ball $B_1^{(0)}(\mathbf{Q}_\infty^\infty(p_n))$ of radius 1 around the vertex $\mathfrak{f}_\infty^{(n)}(0)$. Moreover, still on $\mathcal{E}(n,\alpha)$,

$$d_X([0],t) > 1 \quad \text{whenever } |t| > \alpha.$$

We now define $\mathcal{R}_n \subset \mathbf{Q}_\infty^\infty(p_n) \times \mathcal{T}_X$ by

$$\mathcal{R}_n = \left\{(u, p_X(t)) : u \in A_n, t \in [-\alpha,\alpha] \text{ with } \mathcal{I}(u) = \lfloor t a_n^2 \rfloor\right\}.$$

Letting $\mathbf{E} = (\mathbf{Q}_\infty^\infty(p_n), a_n^{-1}\boldsymbol{d}_{\mathrm{gr}}, \mathfrak{f}_\infty^{(n)}(0))$, $\mathbf{E}' = (\mathcal{T}_X, d_X, [0])$, $r = 1$, we find that given the event $\mathcal{E}(n,\alpha)$, the set $\mathcal{R}_n$ satisfies the requirements of Lemma 10. We are now in the setting of [8, Proof of Theorem 3.5]: All what is left to show is that on $\mathcal{E}(n,\alpha)$, the distortion of $\mathcal{R}_n$ converges to 0 in probability. However, with the same arguments as in the cited proof and using the convergence (17), we obtain

$$\limsup_{n\to\infty} \sup\left\{|\boldsymbol{d}_{\mathrm{gr}}(x,y) - d_X(x',y')| : (x,x'),(y,y') \in \mathcal{R}_n\right\} \leqslant \limsup_{n\to\infty} \frac{5\left(\sup_{A_n} \mathfrak{l}_\infty^{(n)} - \inf_{A_n} \mathfrak{l}_\infty^{(n)}\right)}{a_n}.$$

An appeal to Lemma 7 shows that the right hand side is equal to zero, and the proof of the theorem is completed. □

6 Proofs of the structural properties

6.1 The branching structure behind the UIHPQ_p

In this section, we describe the branching structure of the UIHPQ_p and prove Theorem 4. We will first study a similar mechanism behind Boltzmann quadrangulations Q and Q_σ drawn according to $\mathbb{P}_{g_p,z_p}$ and $\mathbb{P}^\sigma_{g_p}$, respectively (Proposition 6 and Corollary 4), and then pass to the limit $\sigma \to \infty$ using Proposition 2.

To begin with, we follow an idea of [23]: We associate to a (finite) rooted map a tree that describes the branching structure of the boundary of the map. Precisely, for every finite rooted quadrangulation q with a boundary, we define the so-called *scooped-out quadrangulation* $\mathrm{Scoop}(\mathsf{q})$ as follows. We keep only the boundary edges of q and duplicate those edges which lie entirely in the outer face (i.e., whose both sides belong to the outer face). The resulting object is a rooted looptree; see Figure 10.

To a scooped-out quadrangulation we associate its tree of components $\mathrm{Tree}(\mathrm{Scoop}(\mathsf{q}))$ as defined in Section 3.2.4. Following [23], we call this tree, by a slight abuse of terminology, the tree of components of q and use the notation $\mathsf{t} = \mathsf{Tree}(\mathsf{q})$. It is seen that vertices in $\mathsf{t}_\bullet$ have even degree in t, due to the bipartite nature of q.

By gluing the appropriate rooted quadrangulation with a simple boundary into each cycle of $\mathrm{Scoop}(\mathsf{q})$, we recover the quadrangulation q. This provides a bijection

$$\Psi : \mathsf{q} \mapsto (\mathsf{Tree}(\mathsf{q}), (\widehat{\mathsf{q}}_u : u \in V(\mathsf{Tree}(\mathsf{q})_\bullet))$$



Figure 10: A rooted quadrangulation, its boundary and the associated scooped-out quadrangulation.

between, on the one hand, the set $\mathcal{Q}_f$ of finite rooted quadrangulations with a boundary and, on the other hand, the set of plane trees t with vertices at odd height having even degree, together with a collection $(\widehat{\mathsf{q}}_u : u \in V(\mathsf{t}_\bullet))$ of rooted quadrangulations with a simple boundary and respective perimeter $\deg(u)$, for $\deg(u)$ the degree of u in t. We remark that the inverse mapping Ψ^{-1} can be extended to an infinite but locally finite tree together with a collection of quadrangulations with a simple boundary attached to vertices at odd height, yielding in this case an infinite rooted quadrangulation q.

We recall from Section 1.2.5 the definitions of the Boltzmann laws $\mathbb{P}_{g,z}$ and $\mathbb{P}_g^\sigma$, and their analogs with support on quadrangulations with a simple boundary, $\widehat{\mathbb{P}}_{g,z}$ and $\widehat{\mathbb{P}}_g^\sigma$. Their corresponding partition functions are F, F_σ and $\widehat{F}$, $\widehat{F}_\sigma$. We are now interested in the law of the tree of components under $\mathbb{P}_{g,z}$. To begin with, we adapt some enumeration results from [15] to our setting. For every $0 \leqslant p \leqslant 1/2$, recall that $g_p = p(1-p)/3$ and $z_p = (1-p)/4$. Then, (3.15), (3.27) and (5.16) of [15] all together provide the identities

$$F(g_p, z_p) = \frac{2}{3}\frac{3-4p}{1-p}, \quad F_\sigma(g_p) = \frac{(2\sigma)!}{\sigma!(\sigma+2)!}\left(2 + \sigma\frac{1-2p}{1-p}\right)\left(\frac{1}{1-p}\right)^\sigma, \tag{18}$$

for $0 \leqslant p \leqslant 1/2$ and $\sigma \in \mathbb{N}_0$. Moreover, for $\sigma \in \mathbb{N}$ and $0 < p \leqslant 1/2$,

$$\widehat{F}_\sigma(g_p) = \left(\frac{p}{3(1-p)^2}\right)^\sigma \frac{(3\sigma-2)!}{\sigma!(2\sigma-1)!}\left(\frac{3\sigma(1-p)}{p} + 2 - 3\sigma\right), \tag{19}$$

while $\widehat{F}_0(g_p) = 1$. If $p = 0$ and hence $g_p = 0$, then $\widehat{F}_k(0) = \delta_0(k) + \delta_1(k)$ for all $k \in \mathbb{N}_0$. (Indeed, under the maps with no inner faces, the vertex map and the map consisting of one oriented edge are the only maps with a simple boundary.)

We already introduced in Section 2.3 two probability measures $\mu_\circ$ and $\mu_\bullet$ on $\mathbb{N}_0$ given by

$$\mu_\circ(k) = \frac{1}{F(g_p, z_p)}\left(1 - \frac{1}{F(g_p, z_p)}\right)^k, \quad k \in \mathbb{N}_0, \tag{20}$$

$$\mu_\bullet(2k+1) = \frac{1}{F(g_p, z_p) - 1}\left[z_p F^2(g_p, z_p)\right]^{k+1} \widehat{F}_{k+1}(g_p), \quad k \in \mathbb{N}_0, \tag{21}$$

with $\mu_\bullet(k) = 0$ if k even. The tree of components of the scooped-out quadrangulation $\mathrm{Scoop}(Q)$ when Q is drawn according to $\mathbb{P}_{g_p,z_p}$ may now be characterized as follows.

Proposition 6. *Let $0 \leqslant p \leqslant 1/2$, and let Q be distributed according to $\mathbb{P}_{g_p,z_p}$. Then the tree of components $\mathsf{Tree}(Q)$ is a two-type Galton-Watson tree with offspring distribution $(\mu_\circ, \mu_\bullet)$ as given above. Moreover, conditionally on $\mathsf{Tree}(Q)$, the quadrangulations with a simple boundary associated to Q via the bijection Ψ are independent with respective Boltzmann distribution $\widehat{\mathbb{P}}_{g_p}^{\deg(u)}$ for $u \in V(\mathsf{Tree}(Q)_\bullet)$, where $\deg(u)$ denotes the degree of u in $\mathsf{Tree}(Q)$.*

Proof. Note that vertices at even height of $\mathsf{Tree}(Q)$ have an odd number of offspring almost surely. Let t be a finite plane tree satisfying this property. Let also $(\widehat{\mathsf{q}}_u : u \in V(\mathsf{t}_\bullet))$ be a collection of rooted quadrangulations with a simple boundary and respective perimeters $\deg(u)$, and set $\mathsf{q} = \Psi^{-1}(\mathsf{t}, (\widehat{\mathsf{q}}_u : u \in V(\mathsf{t}_\bullet)))$. Then, writing $\Psi_*\mathbb{P}$ for the push-forward measure of $\mathbb{P}$ by Ψ,

$$\Psi_*\mathbb{P}_{g_p,z_p}\left(\mathsf{t}, (\widehat{\mathsf{q}}_u : u \in V(\mathsf{t}_\bullet))\right) = \frac{z_p^{\#\partial\mathsf{q}/2} g_p^{\#\mathrm{F}(\mathsf{q})}}{F(g_p,z_p)} = \frac{1}{F(g_p,z_p)} \prod_{u\in\mathsf{t}_\bullet} z_p^{\deg(u)/2} g_p^{\#\mathrm{F}(\widehat{\mathsf{q}}_u)}.$$

For every $c > 0$, we have

$$1 = \prod_{u\in\mathsf{t}_\circ} c^{k_u} \left(\frac{1}{c}\right)^{\#\mathsf{t}_\bullet} \quad \text{and} \quad \frac{1}{c} = \prod_{u\in\mathsf{t}_\bullet} c^{k_u} \left(\frac{1}{c}\right)^{\#\mathsf{t}_\circ}.$$

Applying the first equality with $c = 1 - 1/F(g_p,z_p)$ and the second one with $c = F(g_p,z_p)$ gives

$$\begin{aligned}\Psi_*\mathbb{P}_{g_p,z_p}\left(\mathsf{t}, (\widehat{\mathsf{q}}_u : u \in \mathsf{t})\right) = \prod_{u\in\mathsf{t}_\circ} \frac{1}{F(g_p,z_p)} \left(1 - \frac{1}{F(g_p,z_p)}\right)^{\deg(u)-1} \\ \times \prod_{u\in\mathsf{t}_\bullet} \frac{1}{F(g_p,z_p)-1} \left(z_p F^2(g_p,z_p)\right)^{\deg(u)/2} \widehat{F}_{\deg(u)/2}(g_p) \prod_{u\in\mathsf{t}_\bullet} \frac{g_p^{\#\mathrm{F}(\widehat{\mathsf{q}}_u)}}{\widehat{F}_{\deg(u)/2}(g_p)},\end{aligned}$$

where we agree that $0/0 = 0$. Therefore,

$$\Psi_*\mathbb{P}_{g_p,z_p}\left(\mathsf{t}, (\widehat{\mathsf{q}}_u : u \in \mathsf{t})\right) = \prod_{u\in\mathsf{t}_\circ} \mu_\circ(k_u) \prod_{u\in\mathsf{t}_\bullet} \mu_\bullet(k_u) \prod_{u\in\mathsf{t}_\bullet} \widehat{\mathbb{P}}_{g_p}^{\deg(u)}(\widehat{\mathsf{q}}_u),$$

which is the expected result. □

Corollary 4. *Let $0 \leqslant p \leqslant 1/2$, $\sigma \in \mathbb{N}$, and let Q be distributed according to $\mathbb{P}_{g_p}^\sigma$. Then the tree of components $\mathsf{Tree}(Q)$ is a two-type Galton-Watson tree with offspring distribution $(\mu_\circ, \mu_\bullet)$ conditioned to have $2\sigma+1$ vertices. Moreover, conditionally on $\mathsf{Tree}(Q)$, the quadrangulations with a simple boundary associated to Q via the bijection Ψ are independent with respective Boltzmann distribution $\widehat{\mathbb{P}}_{g_p}^{\deg(u)}$, for $u \in V(\mathsf{Tree}(Q)_\bullet)$.*

Proof. Observing that $\#V(\mathsf{Tree}(\mathsf{q})) = \#\partial\mathsf{q}+1$ for every rooted quadrangulation q, we obtain

$$\mathbb{P}_{g_p,z_p}(\mathcal{Q}_f^\sigma) = \Psi_*\mathbb{P}_{g_p,z_p}\left(\{\mathsf{t} \in \mathbb{T}_f : \#V(\mathsf{t}) = 2\sigma+1\}\right) = \mathsf{GW}_{\mu_\circ,\mu_\bullet}(\{\mathsf{t} \in \mathbb{T}_f : \#V(\mathsf{t}) = 2\sigma+1\}).$$

Now let t be a finite plane tree with an odd number of offspring at even height, and let $(\widehat{\mathsf{q}}_u : u \in V(\mathsf{t}_\bullet))$ and q be as in the proof of Proposition 6. Then,

$$\Psi_*\mathbb{P}_{g_p}^\sigma\left(\mathsf{t}, (\widehat{\mathsf{q}}_u : u \in \mathsf{t})\right) = \frac{\mathbf{1}_{\{\#\partial\mathsf{q}=2\sigma\}}}{\mathbb{P}_{g_p,z_p}(\mathcal{Q}_f^\sigma)} \prod_{u\in\mathsf{t}_\circ} \mu_\circ(k_u) \prod_{u\in\mathsf{t}_\bullet} \mu_\bullet(k_u) \prod_{u\in\mathsf{t}_\bullet} \widehat{\mathbb{P}}_{g_p}^{\deg(u)}(\widehat{\mathsf{q}}_u),$$

which concludes the proof. □

Lemma 11. *For $0 \leqslant p < 1/2$, the pair $(\mu_\circ, \mu_\bullet)$ is critical and both $\mu_\circ$ and $\mu_\bullet$ have small exponential moments. For $p = 1/2$, the pair $(\mu_\circ, \mu_\bullet)$ is subcritical (and $\mu_\bullet$ has no exponential moment).*

Proof. Recall that $(\mu_\circ, \mu_\bullet)$ is critical if and only if the product of their respective means $m_\circ$ and $m_\bullet$ equals one. Since by (20), $\mu_\circ$ is the geometric law with parameter $1 - 1/F(g_p, z_p)$, we have

$$m_\circ = F(g_p, z_p) - 1.$$

For $m_\bullet$, we let $G_{\mu_\bullet}$ denote the generating function of $\mu_\bullet$. By (21), it follows that

$$G_{\mu_\bullet}(s) = \frac{1}{F(g_p, z_p) - 1} \frac{1}{s} \left(\widehat{F}\left[g_p, z_p F^2(g_p, z_p) s^2\right] - 1 \right), \quad s > 0.$$

Then, Identity (2.8) of [15] ensures that $\widehat{F}(g, zF^2(g, z)) = F(g, z)$ for all non-negative weights g and z. When differentiating this relation with respect to the variable z, we obtain

$$\partial_z \widehat{F}(g, zF^2(g, z)) = \frac{\partial_z F(g, z)}{F^2(g, z) + 2zF(g, z)\partial_z F(g, z)}. \tag{22}$$

Writing

$$\partial_z F(g_p, z_p) = \sum_{\sigma \geqslant 0} \sigma F_\sigma(g_p) z_p^{\sigma - 1},$$

and using the exact expression for $F_\sigma(g_p)$ from (18), we see by means of Stirling's formula that $\partial_z F(g_p, z_p) = \infty$ for $p \in [0, 1/2)$, and $\partial_z F(g_p, z_p) < \infty$ for $p = 1/2$. Thus, for $p \in [0, 1/2)$,

$$\partial_z \widehat{F}(g_p, z_p F^2(g_p, z_p)) = \frac{1}{2 z_p F(g_p, z_p)},$$

whereas if $p = 1/2$, the derivative on the left-hand side in (22) is strictly smaller than the right-hand side for $g = g_p$, $z = z_p$. Finally, applying Identity (2.8) of [15] once again, we get

$$\begin{aligned} m_\bullet =& G'_{\mu_\bullet}(1) \\ =& \frac{1}{F(g_p, z_p) - 1} \left(- \left(\widehat{F}\left[g_p, z_p F^2(g_p, z_p)\right] - 1 \right) + 2 z_p F^2(g_p, z_p) \partial_z \widehat{F}\left[g_p, z_p F^2(g_p, z_p)\right] \right). \end{aligned}$$

As a consequence, $m_\circ m_\bullet = 1$ if $p < 1/2$, and $m_\circ m_\bullet < 1$ if $p = 1/2$. The fact that $\mu_\circ$ has exponential moments is clear. For $\mu_\bullet$, one sees from (19) that the power series

$$\sum_{k \geqslant 0} x^k \widehat{F}_k(g_p)$$

has radius of convergence $\widehat{r}_p = 4(1-p)^2/(9p)$, while (18) ensures that

$$z_p F^2(g_p, z_p) = \frac{(1-4p)^2}{9(1-p)}.$$

Again, for $p \in [0, 1/2)$, $\widehat{r}_p > z_p F^2(g_p, z_p)$, and these quantities are equal for $p = 1/2$. Thus, there exists $s > 1$ such that $G_{\mu_\bullet}(s) < \infty$ if and only if $p < 1/2$, which concludes the proof. □

We are now ready to prove Theorem 4.

Proof of Theorem 4. Fix $0 \leqslant p < 1/2$. Let us denote by Q_∞ the random quadrangulation with an infinite boundary as constructed in the statement of Theorem 4, and let Q_σ be distributed according to $\mathbb{P}^\sigma_{g_p}$. In view of Proposition 2, it is sufficient to prove that in the local sense, as $\sigma \to \infty$,

$$Q_\sigma \xrightarrow{(d)} \mathsf{Q}_\infty. \tag{23}$$

For every real $r \geqslant 1$ and every (finite or infinite) plane tree t, we define $\mathrm{Cut}_r(\mathsf{t})$ as the finite plane tree obtained from pruning all the vertices at a height larger than $2r$ in t. If $\mathsf{q} \in \mathcal{Q}$ is a quadrangulation with a boundary such that $\Psi(\mathsf{q}) = (\mathsf{t}, (\widehat{\mathsf{q}}_u : u \in \mathsf{t}_\bullet))$, we define $\mathrm{Cut}_r(\mathsf{q})$ to be the quadrangulation obtained from gluing the maps $(\widehat{\mathsf{q}}_u : u \in \mathrm{Cut}_r(\mathsf{t})_\bullet)$ in the associated loops of $\mathrm{Loop}(\mathrm{Cut}_r(\mathsf{t}))$. With this definition, we have $B_r(\mathsf{q}) \subset \mathrm{Cut}_r(\mathsf{q})$ for every $r \geqslant 1$, where we recall that $B_r(\mathsf{q})$ stands for the closed ball of radius r around the root in q.

Let $r \geqslant 1$ and $\mathsf{q} \in \mathcal{Q}_f$ such that $\Psi(\mathsf{q}) = (\mathsf{t}, (\widehat{\mathsf{q}}_u : u \in \mathsf{t}_\bullet))$. Using Proposition 6 and Corollary 4, we get

$$\mathbb{P}\left(\mathrm{Cut}_r\left(Q_\sigma\right) = \mathsf{q}\right) = \mathsf{GW}^{(2\sigma+1)}_{\mu_\circ,\mu_\bullet}\left(\mathrm{Cut}_r = \mathsf{t}\right) \prod_{u \in \mathsf{t}_\bullet} \widehat{\mathbb{P}}^{\deg(u)}_{g_p}\left(\widehat{\mathsf{q}}_u\right),$$

where we use the notation $\mathsf{GW}^{(2\sigma+1)}_{\mu_\circ,\mu_\bullet}$ for the $(\mu_\circ, \mu_\bullet)$-Galton-Watson tree conditioned to have $2\sigma + 1$ vertices and interpret Cut_r as the random variable $\mathsf{t} \mapsto \mathrm{Cut}_r(\mathsf{t})$. Applying Proposition 3, we get as $\sigma \to \infty$

$$\mathbb{P}\left(\mathrm{Cut}_r\left(Q_\sigma\right) = \mathsf{q}\right) \longrightarrow \mathsf{GW}^{(\infty)}_{\mu_\circ,\mu_\bullet}\left(\mathrm{Cut}_r = \mathsf{t}\right) \prod_{u \in \mathsf{t}_\bullet} \widehat{\mathbb{P}}^{\deg(u)}_{g_p}\left(\widehat{\mathsf{q}}_u\right) = \mathbb{P}\left(\mathrm{Cut}_r\left(\mathsf{Q}_\infty\right) = \mathsf{q}\right).$$

We proved that for every $r \geqslant 1$, as $\sigma \to \infty$,

$$\mathrm{Cut}_r\left(Q_\sigma\right) \xrightarrow{(d)} \mathrm{Cut}_r\left(\mathsf{Q}_\infty\right).$$

Since $B_r(\mathsf{q}) \subset \mathrm{Cut}_r(\mathsf{q})$ for every $r \geqslant 1$ and $\mathsf{q} \in \mathcal{Q}$, (23) holds and the theorem follows. □

6.2 Recurrence of simple random walk

In this final part, we prove Corollary 2, stating that simple random walk on the UIHPQ_p for $0 \leqslant p < 1/2$ is almost surely recurrent. We will use a criterion from the theory of electrical networks; see, e.g., [35, Chapter 2] for an introduction into these techniques.

Proof of Corollary 2. Fix $0 \leqslant p < 1/2$. We interpret the UIHPQ_p as an electrical network, by equipping each edge with a resistance of strength one. A cutset $\mathcal{C}$ between the root vertex and infinity is a set of edges that separates the root from infinity, in the sense that every infinite self-avoiding path starting from the root has to pass through at least one edge of $\mathcal{C}$. By the criterion of Nash-Williams, cf. [35, (2.13)], it suffices to show that there is a collection $(\mathcal{C}_n, n \in \mathbb{N})$ of disjoint cutsets such that $\sum_{n=1}^\infty (1/\#\mathcal{C}_n) = \infty$ almost surely, i.e., for almost every realization of the UIHPQ_p.

We recall the construction of the UIHPQ_p in terms of the looptree associated to Kesten's two-type tree $\mathcal{T}_\infty = \mathcal{T}_\infty(\mu_\circ, \mu_\bullet)$. Note that the white vertices in $\mathcal{T}_\infty$, i.e., the vertices at even

height, represent vertices in the UIHPQ_p. More precisely, by construction, they form the boundary vertices of the latter. In particular, the white vertices on the spine of $\mathcal{T}_\infty$ are to be found in the UIHPQ_p, and we enumerate them by $v_1, v_2, v_3, \ldots$, such that v_1 is the root vertex, and $d_{\mathrm{gr}}(v_j, v_1) \geqslant d_{\mathrm{gr}}(v_i, v_1)$ for $j \geqslant i$. Now observe that for $i \in \mathbb{N}$, v_i and v_{i+1} lie on the boundary of one common finite-size quadrangulation with a simple boundary, which we denote by $\widehat{\mathfrak{q}}_{v_i}$, in accordance with notation in the proof of Theorem 4.

We define $\mathcal{C}_i$ to be the set of all the edges of $\widehat{\mathfrak{q}}_{v_i}$. Clearly, for each $i \in \mathbb{N}$, $\mathcal{C}_i$ is a cutset between the root vertex and infinity, and for $i \neq j$, $\mathcal{C}_i$ and $\mathcal{C}_j$ are disjoint. The sizes $\#\mathcal{C}_i$, $i \in \mathbb{N}$, are i.i.d. random variables. More specifically, using the construction of the UIHPQ_p in terms of Kesten's looptree, the law of $\#\mathcal{C}_1$ can be described as follows: First, draw a random variable Y according to the size-biased offspring distribution $\bar{\mu}_\bullet$, and then, conditionally on Y, $\#\mathcal{C}_1$ is distributed as the number of edges of a Boltzmann quadrangulation with law $\widehat{\mathbb{P}}_{g_p}^{(Y+1)/2}$, where $g_p = p(1-p)/3$. Obviously, $\#\mathcal{C}_1$ is finite almost surely, implying $\sum_{n=1}^{\infty}(1/\#\mathcal{C}_n) = \infty$ almost surely, and recurrence of the simple symmetric random walk on the UIHPQ_p follows. □

Remark 11. Let us end with a remark concerning the structure of the UIHPQ_p for $p < 1/2$. Note that with probability $\bar{\mu}_\bullet(1) > 0$, a cutset $\mathcal{C}_i$ as constructed in the above proof consists exactly of one edge. By independence and Borel-Cantelli, we thus find with probability one an infinite sequence of such cutsets $\mathcal{C}_{i_1}, \mathcal{C}_{i_2}, \ldots$ consisting of one edge only. In particular, this proves that the UIHPQ_p for $p < 1/2$ admits a decomposition into a sequence of almost surely finite i.i.d. quadrangulations $Q_i(p)$ with a non-simple boundary (whose laws can explicitly be derived from Theorem 4), such that $Q_i(p)$ and $Q_j(p)$ get connected by a single edge if and only if $|i-j| = 1$. This parallels the decomposition of the spaces $\mathbb{H}_\alpha$ for $\alpha < 2/3$ found in [37, Display (2.3)].

Acknowledgments. We warmly thank Grégory Miermont for stimulating discussions and advice.

References

[1] Abraham, C. Rescaled bipartite planar maps converge to the Brownian map. *Ann. Inst. H. Poincaré Probab. Statist.* **52**(2) (2016), 575–595.

[2] Aldous, D. The continuum random tree I. *Ann. Probab.* **19** (1991), 1–28.

[3] Angel, O. Scaling of percolation on infinite planar maps, i. arXiv:0501006.

[4] Angel, O., Curien, N. Percolation on random maps I: half-plane models. *Ann. Inst. H. Poincaré Probab. Statist.* **51**(2) (2015), 405–431.

[5] Angel, O., Ray, G. Classification of half-planar maps. *Ann. Probab.* **43**(3) (2015), 1315–1349.

[6] Angel, O., Ray, G. The half plane UIPT is recurrent. Preprint (2016), arXiv:1601.00410.

[7] Angel, O., Schramm, O. Uniform infinite planar triangulations. *Comm. Math. Phys.* **241**(2-3) (2003), 191–213.

[8] Baur, E., Miermont, G., Ray, G. Classification of scaling limits of uniform quadrangulations with a boundary. Preprint (2016), arXiv:1608.01129.

[9] Baur, E., Miermont, G., Richier, L. Geodesic rays in the Uniform Infinite Half-Planar Quadrangulation return to the boundary. *Lat. Am. J. Probab. Math. Stat.* **13** (2016), 1123–1149.

[10] Benjamini, I., Schramm, O. Recurrence of distributional limits of finite planar graphs. *Electron. J. Probab.* **6**(23) (2001), 13 pp.

[11] Bettinelli, J. Scaling limit of random planar quadrangulations with a boundary. *Ann. Inst. H. Poincaré* **51**(2) (2015), 432–477.

[12] Bettinelli, J., Miermont, G. Compact Brownian surfaces I. Brownian disks. Preprint (2015). arXiv:1507.08776.

[13] Björnberg, J.E., Stefánsson, S. Ö. Recurrence of bipartite planar maps. *Electron. J. Probab.* **19**(31) (2014), 1–40.

[14] Bouttier, J., Di Francesco, P., Guitter, E. Planar maps as labeled mobiles. *Electron. J. Combin.* **11**(1) (2004).

[15] Bouttier, J., Guitter, E. Distance statistics in quadrangulations with a boundary, or with a self-avoiding loop. *J. Phys. A.* **42**(46) (2009), 465208, 44 pp.

[16] Budd, T. The Peeling Process of Infinite Boltzmann Planar Maps. *Electron. J. Combin.* **23**(1), 1–28.

[17] Budzinski, T. The hyperbolic Brownian plane. Preprint (2016). arXiv:1604.06622.

[18] Burago, D., Burago, I., Ivanov, S. *A course in metric geometry.* Graduate Studies in Mathematics (Book 33), AMS (2001).

[19] Caraceni, A., Curien, N. Geometry of the uniform infinite half-planar quadrangulation. Preprint (2015), arXiv:1508.00133.

[20] Curien, N. Peeling random maps. Peccot lecture (2016) Available at http://www.math.u-psud.fr/∼curien/cours/peccot.pdf.

[21] Curien, N. Planar stochastic hyperbolic triangulations. *Probab. Theory Relat. Fields* **165** (2016), 509–540.

[22] Curien, N., Kortchemski, I. Random stable looptrees. *Electron. J. Probab.* **19**(108) (2014), 35 pp.

[23] Curien, N., Kortchemski, I. Percolation on random triangulations and stable looptrees. *Probab. Theory Relat. Fields* **163**(1-2) (2015), 303–337.

[24] Curien, N., Le Gall, J.-F. The Brownian plane. *J. Theor. Probab.* **27**(4) (2014), 1249–1291.

[25] Curien, N., Miermont, G. Uniform infinite planar quadrangulations with a boundary. *Random Struct. Algor.* **47**(1) (2015), 30–58.

[26] Gurel-Gurevich, O., Nachmias, A. Recurrence of planar graph limits. *Ann. of Math.* **177** (2013), 761–781.

[27] Gwynne, E., Miller, J. Scaling limit of the uniform infinite half-plane quadrangulation in the Gromov-Hausdorff-Prokhorov-uniform topology. Preprint (2016), arXiv:1608.00954.

[28] Janson, S. Simply generated trees, conditioned Galton-Watson trees, random allocations and condensation. *Probab. Surveys* **9** (2012), 103–252.

[29] Karatzas, I., Shreve, S. E. *Brownian Motion and Stochastic Calculus.* Graduate Texts in Mathematics (Book 113), Springer, 2nd edition (1991).

[30] Kesten, H. Subdiffusive behavior of random walk on a random cluster. *Ann. Inst. H. Poincaré Probab. Statist.* **22**(4) (1986), 425–487.

[31] Krikun, M. Local structure of random quadrangulations. Preprint (2005), arXiv:0512304.

[32] Le Gall, J.-F. *Spatial branching processes, random snakes and partial differential equations.* Lectures in Mathematics ETH Zürich. Birkhäuser, Basel (1999).

[33] Le Gall, J.-F. Random trees and applications. *Probability Surveys* **2** (2005), 245–311.

[34] Le Gall, J.-F., Miermont, G. *Scaling Limits of Random Trees and Planar Maps.* Clay Mathematics Proceedings **15** (2012).

[35] Lyons, R., Peres, Y. *Probability on trees and networks.* Cambridge University Press, (2016).

[36] Pitman, J. *Combinatorial Stochastic Processes. École d'été de Probabilités de St. Flour.* Lecture Notes in Mathematics **1875**, Springer (2006).

[37] Ray, G. Geometry and percolation on half planar triangulations. *Electron. J. Probab.* **19**(47) (2014), 28 pp.

[38] Richier, L. Universal aspects of critical percolation on random half-planar maps, *Electron. J. Probab.* **20** (2015), 45 pp.

[39] Stephenson, R. Local convergence of large critical multi-type Galton-Watson trees and applications to random maps. Preprint (2014), arXiv:1412.6911. To appear in *J. Theor. Probab.*